


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07D 309/12, C07C 259/06, C07D 213/75, C07C 271/24, C07D 257/04, C07C 259/08, C07D 303/48, 295/18, A61K 31/16, 31/22, 31/395 // C07C 69/675, 69/78, C07D 307/32, 305/12</p>	A1	<p>(11) International Publication Number: <b>WO 98/33788</b></p> <p>(43) International Publication Date: 6 August 1998 (06.08.98)</p>
<p>(21) International Application Number: PCT/EP98/00531</p> <p>(22) International Filing Date: 23 January 1998 (23.01.98)</p> <p>(30) Priority Data: 9702088.7 31 January 1997 (31.01.97) GB</p> <p>(71) Applicant (for all designated States except US): PHARMACIA &amp; UPJOHN S.P.A. [IT/IT]; Via Robert Koch, 1.2, I-20152 Milan (IT).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): ALPEGIANI, Marco [IT/IT]; Via Tolmezzo, 12/5, I-20132 Milan (IT). PAL-LADINO, Massimiliano [IT/IT]; Via Ciro Menotti, 150, I-20025 Legnano (IT). CORIGLI, Riccardo [IT/IT]; Via Morgantini, 31, I-20148 Milan (IT). JABES, Daniela [IT/IT]; Via Costanza, 3, I-20146 Milan (IT). PERRONE, Ettore [IT/IT]; Via Aldo Moro, 44, I-20100 Boffalora Ticino (IT). ABRATE, Francesca [IT/IT]; Via Amedeo d'Aosta, 5, I-20129 Milan (IT). BISSOLINO, Pierluigi [IT/IT]; Via Roma, 36/2, I-27020 San Giorgio Lomellina</p>	<p>(IT). LOMBROSO, Marina [IT/IT]; Via Zanella Giacomo, 48/5, I-20133 Milan (IT).</p> <p>(81) Designated States: AU, BR, CA, CN, HU, IL, JP, KR, MX, NO, NZ, PL, UA, US, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p><b>Published</b> With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	
<p>(54) Title: MATRIX METALLOPROTEINASE INHIBITORS</p> <p>(57) Abstract</p> <p>A compound of formula (I), wherein W is -NHOH or -OH, R<sub>1</sub> is either free or protected hydroxymethyl or mercaptomethyl or derivatives thereof, R<sub>2</sub> is free or protected hydroxy, R<sub>3</sub> and R<sub>4</sub> is an organic group, R<sub>5</sub> is hydrogen or methyl, or R<sub>4</sub> and R<sub>5</sub>, together with the nitrogen atom to which they are attached, constitute an azaheterocyclyl group, and the solvates, hydrates and pharmaceutically acceptable salts thereof, can inhibit matrix metalloproteinases and the release of tumour necrosis factor (TNF). Processes for producing the compound, intermediates involved in the processes, and pharmaceutical compositions containing the compound are also described.</p> <div style="text-align: right;">  <p>(I)</p> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## MATRIX METALLOPROTEINASE INHIBITORS

The present invention relates to new inhibitors of  
5 matrix metalloproteinases (hereinafter MMPs), to a process  
for their preparation, to pharmaceutical compositions  
containing them, and to the use of such compounds in the  
prevention, control and treatment of diseases in which the  
proteolytic action of MMPs is involved. Furthermore, since  
10 some of the compounds herein described potently inhibit the  
release of tumor necrosis factor-alpha (hereinafter TNF) from  
cells, another object of the present invention is the use of  
pharmaceutical compositions containing said compounds for the  
treatment or prophylaxis of inflammatory, immunological or  
15 infectious diseases promoted by the release of such cytokine.

Low molecular weight compounds able to inhibit one or  
more of the matrix metalloproteinases, in particular  
stromelysin-1 (MMP-3; EC 3.4.24.17), gelatinase A (MMP-2; EC  
3.4.24.24), gelatinase B (MMP-9; EC 3.4.24.35), neutrophil  
20 collagenase (MMP-8; EC 3.4.24.34), interstitial collagenase  
(MMP-1; EC 3.4.27.7), matrilysin (MMP-7; EC 3.4.24.23), and  
collagenase-3 (MMP-13) are currently considered as promising  
therapeutic agents in degenerative, tumoral and autoimmune  
pathologies (e.g., P.D. Brown: "Matrix metalloproteinase  
25 inhibitors: A new class of anticancer agent", Curr. Opin.  
Invest. Drugs, 2:617-626, 1993; A. Krantz: "Proteinases in  
Inflammation", Annu. Rep. Med. Chem. 28:187-195, 1993). Many  
of such compounds described hitherto are peptide derivatives  
or pseudopeptides, bearing analogies to recognized peptide  
30 substrates of these enzymes, and characterized in addition by  
a functional group capable of binding the Zn (II) atom

present in the catalytic site of said enzymes. Known classes of MMP inhibitors include those in which the Zn binding group is a hydroxamic acid, which is part of a (substituted) succinic moiety, in particular a succinic amide represented by the general formula (A)




wherein  $R_a$ ,  $R_b$ ,  $R_c$ , and  $R_d$  are hydrogen atoms or appropriate substituents (e.g., N.R.A. Beeley et al., "Inhibitors of matrix metalloproteinases (MMP's)", Curr. Opin. Ther. Patents 4:7-16, 1994; J.R. Porter et al., "Recent developments in matrix metalloproteinase inhibitors", Exp. Opin. Ther. Patents 5:1287-1296, 1995; J.R. Morphy et al., "Matrix metalloproteinase inhibitors: Current status", Curr. Med. Chem. 2:743-762, 1995; R.P. Beckett et al., "Recent advances in matrix metalloproteinase research", DDT 1:16-26, 1996).

Further, it is now recognized that some compounds of the same general formula (A) may be able to inhibit the release of TNF from the cell membrane anchored precursor, pro-TNF (e.g., G.M. McGeehan et al., "Regulation of tumour necrosis factor-alpha processing by a metalloproteinase inhibitor", Nature 370:558-561, 1994).

Although MMPs have been recognized as drug targets for at least 20 years, and MMP inhibitors encompassed by the general formula (A) have been disclosed since 1986 or before (e.g., see J.P. Dickens et al., U.S. Patent 4,599,361), no drug of this type has arrived at the market yet. This is not because of questions about the therapeutic potential of MMP

inhibitors, but because of problems of the "first generation" compounds, such as inhibitor potency, aqueous solubility, metabolic stability, and other desirable properties, oral bioavailability in particular (e.g., J.R. Porter, reference above; J. Hodgson, "Remodelling MMPs", Biotechnology 13:554-557, 1995). For example, it is well known that most "first generation" hydroxamate MMP inhibitors are rapidly glucuronidated, oxidized to the carboxylic acid, and excreted in the bile (e.g., see J. Singh et al., Bioorg. Med. chem. Lett. 5:337-342, 1995, and reference above). Thus, there is a strong need for better and diversified molecules, especially as far as the properties referred to above are concerned. The present invention relates to a new class of carboxylic or hydroxamic acid derivatives, which are characterised by the presence of a 1,2-glycol, or a derivative thereof, next to the carbonyl group.

In particular, the present invention provides a compound of formula (I)



20


wherein

W is -NHOH or -OH;


R<sub>1</sub> is hydroxymethyl or a hydroxymethyl derivative which is an ether, an ester, a carbonate or a carbamate; or

25 R<sub>1</sub> is mercaptomethyl or a mercaptomethyl derivative which is a sulfide (i.e., a thioether), a sulfoxide, a sulfone or a thioester;

$R_2$  is hydroxy or a protected derivate thereof; or  
 $R_1$  and  $R_2$ , taken together with the carbon atom to which they  
 are attached, represent an optionally substituted  
 cyclopropane or oxirane, 1,3-dioxolane or 2-oxo-1,3-dioxolane  
 5 ring of the following formulae (B1)-(B4):



$R_3$  is a group  $-A^I-X-(CH_2)_n-A$ , wherein A is  $C_1-C_{10}$  alkyl,  $C_2-C_{10}$   
 alkenyl,  $C_3-C_7$  cycloalkyl, aryl, or heterocyclyl, the said  
 10 alkyl, alkenyl, cycloalkyl, aryl and heterocyclyl groups  
 being unsubstituted or substituted; n is either zero or an  
 integer from 1 to 5; -X- is either a direct bond or a group -  
 O-, -S-, -SO-, -SO<sub>2</sub>-, -SO<sub>2</sub>NH-, -CO-, -CONH-, -NHCO-, -OCONH-,  
 NHCONH or -NHSO<sub>2</sub>-, and  $-A^I$  is  $C_1-C_{10}$  alkylene,  $C_2-C_6$   
 15 alkenylene, or phenylene;  
 $R_4$  is either a group of formula (C):




wherein  $R_6$  is hydrogen or the side chain of a natural or non-  
 20 natural alpha-amino acid, and  $R_7$  is amino or a group -NH-A, -  
 NH-CH<sub>2</sub>-A or -NH-CH<sub>2</sub>CH<sub>2</sub>-A wherein A is as defined above; or


5

$R_4$  is a group A as defined above or a group  $-A^1-X-A$  wherein A,  $-X-$  and  $-A^1-$  are as defined above; or

$R_4$  is a group of formula (D):



- wherein  $R_8$  is methyl, ethyl, phenyl, pyridyl, isopropyl, sec-butyl, tert-butyl, adamantyl, cyclopentyl, cyclohexyl, cyclohexylmethyl, phenylmethyl, 3,4-(methylenedioxy)phenyl, piperonyl, pyridylmethyl, thienylmethyl, 3-indolylmethyl, 3-(N-methyl)indolylmethyl, 1,1-diphenylmethyl, naphthyl, naphthylmethyl, quinolylmethyl, isoquinolylmethyl, thiazolyl, thiazolylmethyl, imidazolyl, imidazolylmethyl, or a derivative thereof optionally substituted, or  $R_8$  is  $-C(CH_3)_2SCH_3$  or a sulfoxide or sulfone thereof,  $-C(CH_3)_2OCH_3$ ,  $-CH_2CH_2CH_2-OCH_3$ , or  $-CH(CH_3)OH$  or a tert-butyl ether thereof;  $R_9$  is either hydrogen or a group selected from methyl, ethyl, phenyl, 3,4-(methylenedioxy)phenyl, piperonyl, pyridyl, benzimidazolyl and 4-tetrazolyl, which group is optionally substituted; or  $R_8$  and  $R_9$ , taken together with the nitrogen atom to which they are attached, constitute a dihydrocarbostyryl ring (D'):



wherein the nitrogen atom and the phenyl ring may be optionally substituted;

$R_5$  is hydrogen or methyl; or

$R_4$  and  $R_5$ , taken together with the nitrogen atom to which they are attached, form an optionally substituted azaheterocyclyl ring; and the solvates, hydrates and pharmaceutically acceptable salts thereof.

As used herein the term "alkyl" refers to a straight or branched chain alkyl moiety having from 1 to 10 carbon atoms, including for example, methyl, ethyl, propyl, isopropyl, n-butyl, i-butyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, n-hexyl, n-heptyl and n-octyl. The term "alkylene" refers to an alkyl group as defined above, but connected to the rest of the molecule by two covalent bonds, either at the same or at different carbon atoms, including for example methylene ( $-\text{CH}_2-$ ), 1,2-ethylene ( $-\text{CH}_2-\text{CH}_2-$ ) and 1,1-ethylene ( $-\text{CH}(\text{CH}_3)-$ ). The term "alkenyl" as used herein refers to a straight or branched chain alkenyl moiety having from 2 to 10 carbon atoms and having in addition one double bond of either E or Z stereochemistry where applicable. Examples of alkenyl groups include: vinyl, allyl, metallyl, butenyl and crotyl. The term "alkenylene" refers to an alkenyl group as defined above, but connected to the rest of the molecule by two covalent bonds, either at the same or at different carbon atoms, including for example 1,2-ethenylene ( $-\text{CH}=\text{CH}-$ ) and 1,1-ethenylene ( $-\text{C}(=\text{CH}_2)-$ ) and 1,3-propenylene ( $-\text{CH}=\text{CH}-\text{CH}_2-$ ).

The term "aryl" as used herein refers to a monocyclic or bicyclic aromatic hydrocarbon group of 6 to 10 carbon atoms, such as phenyl, naphthyl, indanyl; furthermore, "aryl" as used herein may refer to a biphenyl group ( $-\text{C}_6\text{H}_4-\text{C}_6\text{H}_5$ ). The term "phenylene" refers to a phenyl group connected to the rest of



the molecule by two covalent bonds, i.e. 1,2-C<sub>6</sub>H<sub>4</sub> (ortho-phenylene), 1,3-C<sub>6</sub>H<sub>4</sub> (meta-phenylene), and 1,4-C<sub>6</sub>H<sub>4</sub> (para-phenylene).

The term "cycloalkyl" as used herein refers to a  
5 saturated carbocyclic group of 3 to 7 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

The term "heterocyclyl" as used herein refers to a 3-  
to 7-membered, saturated or unsaturated heterocyclyl ring,  
10 containing at least one heteroatom selected from O, S and N, wherein any ring nitrogen may be oxidized as an N-oxide, any ring carbon may be oxidized as a carbonyl, and any ring sulfur may be oxidized as a sulfoxide or sulfone; and wherein said heterocyclyl ring may be optionally fused to a second 5- or  
15 6-membered, saturated or unsaturated heterocyclyl ring, or to a C<sub>3</sub>-C<sub>7</sub> cycloalkyl ring, or to a benzene or naphthalene ring.

Examples of heterocyclyl groups are pyrrolyl,  
pyrrolidinyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl,  
oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazolyl,  
20 thienyl, tetrahydrothienyl, furyl, tetrahydrofuryl, aziridinyl, oxiranyl, azetidiny, succinimido, pyridyl, piperidinyl, pyrazinyl, piperazinyl, pyridazinyl, hexahydropyridazinyl, pyrimidinyl, pyranyl, tetrahydropyranyl, benzothienyl, benzothiazolyl, benzoxazolyl, isobenzofuranyl, benzofuranyl, benzimidazolyl, indazolyl, chromenyl, indolyl,  
25 oxindolyl, phthalimido, 1-oxo-2-isoindolyl, quinolyl, isoquinolyl, tetrahydroisoquinolyl, indoliziny, isoindolyl, 2-oxoisoindolyl, 1,2-(methylenedioxy)phenyl, quinuclidinyl, hydantoinyl, saccharinyl, cinnolinyl, purinyl, morpholinyl,  
30 thiomorpholinyl, dioxanyl, dithianyl and azepinyl.

The term "azaheterocyclyl" as used herein includes any of the heterocyclyl groups as defined above which contain at least one nitrogen atom and which are linked to the rest of the molecule by a nitrogen atom.

5       The term "side chain of a naturally occurring  $\alpha$ -amino acid" encompasses the side chains of alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine,  
10   tryptophan, tyrosine, valine, and penicillamine, and derivatives thereof wherein any -OH, -SH, -NH, and -NH<sub>2</sub> may be alkylated by benzyl or C<sub>1</sub>-C<sub>6</sub> linear or branched alkyl groups, or acylated by benzoyl or C<sub>2</sub>-C<sub>7</sub> linear or branched alkanoyl groups.

15       The term "side chain of a non-natural  $\alpha$ -amino acid" encompasses the side chain of known  $\alpha$ -amino acids not belonging to the category of "naturally occurring  $\alpha$ -amino acids", such as  $\alpha$ -amino-*n*-butyric acid,  $\alpha$ -amino-*n*-pentanoic acid,  $\alpha$ -amino-*n*-hexanoic acid,  $\alpha$ -amino-neohexanoic acid,  $\alpha$ -  
20   amino-neoheptanoic acid, S-methyl penicillamine sulfoxides and sulfone, tert-butylglycine (alias tert-leucine), (2-methoxy-2-propyl)glycine, (3-methoxy-1-propyl)glycine, phenylglycine, (diphenylmethyl)-glycine, quinolylalanine, isoquinolylalanine, pyridylglycine, (2-pyridyl)alanine, (3-  
25   pyridyl)alanine, (4-pyridyl)alanine, adamantylglycine, cyclohexylalanine, cyclohexylglycine, homophenylalanine, naphthylalanine, thienylalanine, homocysteine, homoserine, alloseleucine, allothreonine and ornithine. Included in this definition, as used herein, are also side chains of natural  
30    $\alpha$ -amino acids, as defined above, which are substituted at any carbon atom by one or more substituents selected from chloro,

fluoro, hydroxy, methoxy and trifluoromethyl, such as, for example, 3,4-dihydroxyphenylalanine, 5-hydroxylysine and 4-fluorophenylalanine.

In the above definition of  $R_1$ , a "hydroxymethyl derivative" is an ether, an ester, a carbonate or a carbamate. More particularly,  $R_1$  can be  $-\text{CH}_2-\text{O}-(\text{CH}_2)_n-\text{A}$ ,  $-\text{CH}_2-\text{O}-\text{C}(\text{O})-(\text{CH}_2)_n-\text{A}$ ,  $-\text{CH}_2-\text{O}-\text{C}(\text{O})-\text{O}-(\text{CH}_2)_n-\text{A}$ ,  $-\text{CH}_2-\text{O}-\text{C}(\text{O})-\text{NH}-(\text{CH}_2)_n-\text{A}$ , or  $-\text{CH}_2-\text{O}-\text{C}(\text{O})-\text{N}(\text{A})-(\text{CH}_2)_n-\text{A}'$ , wherein  $n$  and  $\text{A}$  are as defined above, and  $\text{A}'$ , being the same or different, is a group  $\text{A}$  as defined above, and wherein the residue  $-\text{N}(\text{A})-(\text{CH}_2)_n-\text{A}'$  may constitute, in addition, a five- or six-membered azaheterocyclic ring selected from morpholine, pyrrolidine, piperidine, piperazine and hexahydropyridazine, optionally substituted by methyl, phenyl, benzyl, chloro, fluoro, methoxy and carbamoyl. Ether derivatives of such a hydroxymethyl group include the ether derivatives commonly used as hydroxy-protecting groups, such as methoxymethyl ethers and tetrahydropyranyl ethers.

In the above definition of  $R_1$ , a "mercaptomethyl derivative" is a sulfide, a sulfoxide, a sulfone or a thioester. More particularly  $R_1$  can be  $-\text{CH}_2-\text{S}-(\text{CH}_2)_n-\text{A}$ ,  $-\text{CH}_2-\text{S}(\text{O})-(\text{CH}_2)_n-\text{A}$ ,  $-\text{CH}_2-\text{S}(\text{O})_2-(\text{CH}_2)_n-\text{A}$ , or  $-\text{CH}_2-\text{S}-\text{C}(\text{O})-(\text{CH}_2)_n-\text{A}$ , wherein  $\text{A}$  and  $n$  are as defined above.

When  $R_1$  and  $R_2$ , taken together with the carbon atom to which they are attached, form a substituted cyclopropane or oxirane ring, such rings are preferably substituted by one or two substituents selected from  $\text{C}_1-\text{C}_3$  alkyl, phenyl and benzyl. When  $R_1$  and  $R_2$ , taken together with the carbon atom to which they are attached, form a substituted 1,3-dioxolane ring, this ring is preferably substituted by one or two substituents selected from  $\text{C}_1-\text{C}_3$  alkyl,  $\text{C}_1-\text{C}_3$  alkoxy, phenyl and benzyl.

Substituents which may be present in the methyl, alkyl, alkenyl, aryl, cycloalkyl, heterocyclyl and azaheterocyclyl groups in any of the above definitions of  $R_1$ - $R_7$ , include the following:

- 5 - halo (i.e., fluoro, bromo, chloro or iodo);
- hydroxy;
- nitro;
- azido;
- mercapto (i.e., -SH), and acetyl or phenylacetyl esters
- 10 thereof (i.e., -SCOCH<sub>3</sub> and -SCOCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>);
- amino (i.e., -NH<sub>2</sub> or -NHR<sup>I</sup> or -NR<sup>I</sup>R<sup>II</sup>, wherein R<sup>I</sup> and R<sup>II</sup>, which are the same or different, are straight or branched C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, biphenyl (i.e., -C<sub>6</sub>H<sub>4</sub>-C<sub>6</sub>H<sub>5</sub>), or benzyl groups, optionally substituted by hydroxy, methoxy, methyl,
- 15 amino, methylamino, dimethylamino, chloro or fluoro; or R<sup>I</sup> and R<sup>II</sup> taken together with the nitrogen atom to which they are attached form a heterocyclic ring such as morpholino, pyrrolidino, piperidino, pyperazino or N-methylpyperazino;
- guanidino, i.e., -NHC(=NH)NH<sub>2</sub>;
- 20 - formyl (i.e. -CHO);
- cyano;
- carboxy (i.e. -COOH), or esters thereof (i.e., -COOR<sup>I</sup>), or amides thereof (i.e., -CONH<sub>2</sub>, -CONHR<sup>I</sup> or -CONHR<sup>I</sup>R<sup>II</sup>), wherein R<sup>I</sup> and R<sup>II</sup> are as defined above, and including
- 25 morpholino-amides, pyrrolidino-amides, and carboxymethylamides -CONHCH<sub>2</sub>COOH;
- sulfo (i.e., -SO<sub>3</sub>H);
- acyl, i.e., -C(O)R<sup>I</sup>, wherein R<sup>I</sup> is as defined above, including monofluoroacetyl, difluoroacetyl, trifluoroacetyl;
- 30 - carbamoyloxy (i.e., -OCONH<sub>2</sub>) and N-methylcarbamoyloxy;

- 11
- acyloxy, i.e.,  $-\text{OC}(\text{O})\text{R}^{\text{I}}$  wherein  $\text{R}^{\text{I}}$  is as defined above, or formyloxy;
  - acylamino, i.e.,  $-\text{NHC}(\text{O})\text{R}^{\text{I}}$ , or  $-\text{NHC}(\text{O})\text{OR}^{\text{I}}$ , wherein  $\text{R}^{\text{I}}$  is as defined above or is a group  $-(\text{CH}_2)_t\text{COOH}$  where  $t$  is 1, 2 or 3;
  - ureido, i.e.,  $-\text{NH}(\text{CO})\text{NH}_2$ ,  $-\text{NH}(\text{CO})\text{NHR}^{\text{I}}$ ,  $-\text{NH}(\text{CO})\text{NR}^{\text{I}}\text{R}^{\text{II}}$ , wherein  $\text{R}^{\text{I}}$  and  $\text{R}^{\text{II}}$  are as defined above, including  $-\text{NH}(\text{CO})-(4\text{-morpholino})$ ,  $-\text{NH}(\text{CO})-(1\text{-pyrrolidino})$ ,  $-\text{NH}(\text{CO})-(1\text{-piperazino})$ ,  $-\text{NH}(\text{CO})-(4\text{-methyl-1-piperazino})$ ;
  - sulfonamido, i.e.,  $-\text{NHSO}_2\text{R}^{\text{I}}$  wherein  $\text{R}^{\text{I}}$  is as defined above;
  - a group  $-(\text{CH}_2)_t\text{COOH}$ , and esters and amides thereof, i.e.,  $-(\text{CH}_2)_t\text{COOR}^{\text{I}}$  and  $-(\text{CH}_2)_t\text{CONH}_2$ ,  $-(\text{CH}_2)_t\text{CONHR}^{\text{I}}$ ,  $-(\text{CH}_2)_t\text{CONR}^{\text{I}}\text{R}^{\text{II}}$ , wherein  $t$ ,  $\text{R}^{\text{I}}$  and  $\text{R}^{\text{II}}$  are as defined above;
  - a group  $-\text{NH}(\text{SO}_2)\text{NH}_2$ ,  $-\text{NH}(\text{SO}_2)\text{NHR}^{\text{I}}$ ,  $-\text{NH}(\text{SO}_2)\text{NR}^{\text{I}}\text{R}^{\text{II}}$ , wherein  $\text{R}^{\text{I}}$  and  $\text{R}^{\text{II}}$  are as defined above, including  $-\text{NH}(\text{SO}_2)-(4\text{-morpholino})$ ,  $-\text{NH}(\text{SO}_2)-(1\text{-pyrrolidino})$ ,  $-\text{NH}(\text{SO}_2)-(1\text{-piperazino})$ ,  $-\text{NH}(\text{SO}_2)-(4\text{-methyl-1-piperazino})$ ;
  - a group  $-\text{OC}(\text{O})\text{OR}^{\text{I}}$ , wherein  $\text{R}^{\text{I}}$  is as defined above;
  - a group  $-\text{OR}^{\text{I}}$ , wherein  $\text{R}^{\text{I}}$  is as defined above, including  $-\text{OCH}_2\text{COOH}$ ;
  - a group  $-\text{SR}^{\text{I}}$ , wherein  $\text{R}^{\text{I}}$  is as defined above, including  $-\text{SCH}_2\text{COOH}$ ;
  - a group  $-\text{S}(\text{O})\text{R}^{\text{I}}$ , wherein  $\text{R}^{\text{I}}$  is as defined above;
  - a group  $-\text{S}(\text{O}_2)\text{R}^{\text{I}}$ , wherein  $\text{R}^{\text{I}}$  is as defined above;
  - a group  $-\text{SO}_2\text{NH}_2$ ,  $-\text{SO}_2\text{NHR}^{\text{I}}$ , or  $-\text{SO}_2\text{NR}^{\text{I}}\text{R}^{\text{II}}$ , wherein  $\text{R}^{\text{I}}$  and  $\text{R}^{\text{II}}$  are as defined above;
  - $\text{C}_1$ - $\text{C}_6$  alkyl or  $\text{C}_2$ - $\text{C}_6$  alkenyl;
  - $\text{C}_3$ - $\text{C}_7$  cycloalkyl;
  - substituted methyl selected from chloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, aminomethyl, N,N-

dimethylaminomethyl, azidomethyl, cyanomethyl, carboxymethyl, sulfomethyl, carbamoylmethyl, carbamoyloxymethyl, hydroxymethyl, methoxycarbonylmethyl, ethoxycarbonylmethyl, tert-butoxycarbonylmethyl and guanidinomethyl.

5       When present, carboxy, hydroxy, mercapto and amino groups may be either free or in a protected form. Protected forms of said groups are any of those generally known in the art, as described, for example, by T.W. Greene in "Protective Groups in Organic Chemistry", Wiley Interscience. Preferably, 10 carboxy groups are protected as esters thereof, in particular methyl, ethyl, tert-butyl, benzyl, and 4-nitrobenzyl esters. Preferably, hydroxy groups are protected as ethers or esters thereof, in particular methoxymethyl ethers, tetrahydropyranyl ethers, benzyl ethers, acetates or benzoates. Preferably, 15 mercapto groups are protected as thioethers or thioesters, in particular tert-butyl thioethers, thioacetates or thiobenzoates. Preferably, amino groups are protected as carbamates, e.g. tert-butoxycarbonyl derivatives, or as amides, e.g. acetamides and benzamides.

20       The present invention provides the salts of those compounds of formula (I) that have salt-forming groups, especially the salts of the compounds having a carboxylic group, a N-hydroxycarbamoyl group, and a sulfo group, or the salts of the compounds having a basic group, especially an 25 amino or guanidino group. The salts are especially physiologically tolerable salts, for example alkali metal and alkaline earth metal salts (e.g. sodium, potassium, lithium, calcium and magnesium salts), ammonium salts and salts with an appropriate organic amine or amino acid (e.g. arginine, 30 procaine salts), and the addition salts formed with suitable inorganic acids (e.g. hydrochlorides, hydrobromides,


sulfates, phosphates) or carboxylic and sulfonic organic acids (e.g. acetates, trifluoroacetates, citrates, succinates, malonates, lactates, tartrates, fumarates, maleates, methanesulfonates, *p*-toluenesulfonates). Some compounds of formula (I) which contain a carboxylate and an ammonium group may exist as zwitterions; such salts are also part of the present invention.

Furthermore, hydrates, solvates of compounds of formula (I), and physiologically hydrolyzable derivatives (i.e., prodrugs) of compounds of formula (I) are included within the scope of the present invention.

The present invention encompasses all the possible stereoisomers (e.g. diastereoisomers, epimers, geometrical isomers) of the compounds of formula (I), as well as their racemic or optically active mixtures related to the  $R_1$ - $R_5$  substituents.

The present invention also provides a pharmaceutical compositions which comprises, as active ingredient, an amide derivative of formula (I) or a solvate hydrate or pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier, diluent or excipient.

Preferred compounds of the present invention are represented by the formula (I')



25

wherein:

W is -NHOH or -OH;


- $R_1$  is hydroxymethyl or the ether, ester, carbonate and carbamate derivatives thereof, as defined above; or
- $R_1$  is a mercaptomethyl or the sulfide, sulfone and thioester derivatives thereof, as defined above;
- 5  $R_2$  is hydroxy or a protected derivative thereof; or
- $R_1$  and  $R_2$ , taken together with the carbon atom to which they are attached, form a cyclopropane or oxirane ring, optionally substituted by phenyl or benzyl;
- $R_3$  is  $-\text{CH}_2\text{-alkyl}$ ,  $-\text{CH}_2\text{-cycloalkyl}$ ,  $-(\text{CH}_2)_n\text{-O-alkyl}$ ,  $-(\text{CH}_2)_n\text{-O-}$
- 10  $\text{cycloalkyl}$ ,  $-(\text{CH}_2)_n\text{-O-}(\text{CH}_2)_{n'}\text{-aryl}$ ,  $-(\text{CH}_2)_n\text{-O-}(\text{CH}_2)_{n'}\text{-}$
- $\text{heterocyclyl}$ ,  $-(\text{CH}_2)_n\text{-S-alkyl}$ ,  $-(\text{CH}_2)_n\text{-S-cycloalkyl}$ ,  $-(\text{CH}_2)_n\text{-S-}$
- $(\text{CH}_2)_{n'}\text{-aryl}$ ,  $-(\text{CH}_2)_n\text{-S-}(\text{CH}_2)_{n'}\text{-heterocyclyl}$ ,  $-(\text{CH}_2)_n\text{-S(O)-}$
- $\text{alkyl}$ ,  $-(\text{CH}_2)_n\text{-S(O)-cycloalkyl}$ ,  $-(\text{CH}_2)_n\text{-SO}_2\text{-alkyl}$ ,  $-(\text{CH}_2)_n\text{-SO}_2\text{-}$
- $\text{cycloalkyl}$ ,  $-(\text{CH}_2)_n\text{-SO}_2\text{-}(\text{CH}_2)_{n'}\text{-aryl}$ ,  $-(\text{CH}_2)_n\text{-SO}_2\text{-}(\text{CH}_2)_{n'}\text{-}$
- 15  $\text{heterocyclyl}$ ,  $-(\text{CH}_2)_n\text{-SO}_2\text{NH-alkyl}$ ,  $-(\text{CH}_2)_n\text{-SO}_2\text{NH-cycloalkyl}$ ,  $-(\text{CH}_2)_n\text{-SO}_2\text{NH-}(\text{CH}_2)_{n'}\text{-aryl}$ ,  $-(\text{CH}_2)_n\text{-SO}_2\text{NH-}(\text{CH}_2)_{n'}\text{-heterocyclyl}$ ,  $-(\text{CH}_2)_n\text{-CO-alkyl}$ ,  $-(\text{CH}_2)_n\text{-CO-cycloalkyl}$ ,  $-(\text{CH}_2)_n\text{-CO-}(\text{CH}_2)_{n'}\text{-aryl}$ ,  $-(\text{CH}_2)_n\text{-CO-}(\text{CH}_2)_{n'}\text{-heterocyclyl}$ ,  $-(\text{CH}_2)_n\text{-CONH-alkyl}$ ,  $-(\text{CH}_2)_n\text{-CONH-cycloalkyl}$ ,  $-(\text{CH}_2)_n\text{-CONH-}(\text{CH}_2)_{n'}\text{-aryl}$ ,  $-(\text{CH}_2)_n\text{-CONH-}$
- 20  $(\text{CH}_2)_{n'}\text{-heterocyclyl}$ ,  $-(\text{CH}_2)_n\text{-NHCO-alkyl}$ ,  $-(\text{CH}_2)_n\text{-NHCO-cycloalkyl}$ ,  $-(\text{CH}_2)_n\text{-NHCO-}(\text{CH}_2)_{n'}\text{-aryl}$ ,  $-(\text{CH}_2)_n\text{-NHCO-}(\text{CH}_2)_{n'}\text{-heterocyclyl}$ ,  $-(\text{CH}_2)_n\text{-NHSO}_2\text{-alkyl}$ ,  $-(\text{CH}_2)_n\text{-NHSO}_2\text{-cycloalkyl}$ ,  $-(\text{CH}_2)_n\text{-NHSO}_2\text{-}(\text{CH}_2)_{n'}\text{-aryl}$  or  $-(\text{CH}_2)_n\text{-NHSO}_2\text{-}(\text{CH}_2)_{n'}\text{-heterocyclyl}$ , wherein the said alkyl, cycloalkyl, aryl and heterocyclyl
- 25 groups are unsubstituted or substituted by one to three substituents selected from chloro, fluoro, hydroxy, methoxy, and  $\text{C}_1\text{-C}_4$  alkyl; and  $n$  and  $n'$ , being the same or different, are zero or an integer from 1 to 5; more preferably,  $R_3$  is selected from the group consisting of  $R_3$  is isobutyl,
- 30 cyclopentylmethyl, 3-(4-chlorophenyl)propyl, 3-(4-methoxyphenyl)propyl, 3-(4-biphenyl)propyl, 3-[4-(4-



15  
 fluorophenyl)phenyl]propyl, 3-[4-(4-  
 chlorophenyl)phenyl]propyl, 3-[4-(4-  
 methoxyphenyl)phenyl]propyl, 3-(4-phenoxyphenyl)propyl, 3-(4-  
 pyridoxyphenyl)propyl, (4-methoxybenzene)sulfonylmethyl, and  
 5 (4-butoxybenzene)sulfonyl-methyl;  
 and  $R_4$ ,  $R_5$  are as defined above.

A first group of particularly preferred compound are  
 those of the formula (I'), wherein  $W$ ,  $R_1$ ,  $R_2$  and  $R_3$  are as  
 defined above, and  $R_4$  is a group of formula (C):

10




wherein  $R_6$  is phenyl, pyridyl, isopropyl, sec-butyl, tert-  
 butyl, adamantyl, cyclopentyl, cyclohexyl, cyclohexylmethyl,  
 phenylmethyl, -pyridylmethyl, 3-pyridylmethyl, 4-  
 15 pyridylmethyl, 2-thienylmethyl, 3-indolylmethyl, 3-(N-  
 methyl)indolylmethyl, 1,1-diphenylmethyl, 1-naphthyl, 2-  
 naphthyl, 1-naphthylmethyl, 2-naphthylmethyl, quinolylmethyl  
 or isoquinolylmethyl, or a derivative thereof substituted by  
 one to three substituents selected from chloro, fluoro,  
 20 hydroxy, methoxy, carbamoyl and  $C_1$ - $C_4$  alkyl; or  $R_6$  is -  
 $C(CH_3)_2SCH_3$  or a sulfoxide or sulfone thereof,  $-C(CH_3)_2OCH_3$ , -  
 $CH_2CH_2CH_2-OCH_3$ , or  $-CH(CH_3)OH$  or the tert-butyl ether thereof;  
 and wherein  $R_7$  is  $-NH_2$  or a group  $-NH-A-$ ,  $NH-CH_2-A$  or  $NH-$   
 $CH_2CH_2-A$ , wherein  $A$ , being as defined above, is selected from  
 25 methyl, isopropyl, tert-butyl, phenyl, 4-methoxyphenyl, 4-  
 fluorophenyl, 4-(aminosulfonyl)phenyl, 4-(dimethylamino-  
 sulfonylmethyl)phenyl, (3,4-methylenedioxy)phenyl, 2-pyridyl,

16

3-pyridyl, 4-pyridyl, 4-piperidyl, 2-thiazolyl, 1-naphthyl,  
 2-naphthyl, 3-quinolyl, 5-quinolyl, 3-isoquinolinyl, 5-  
 isoquinolyl, 3-quinuclidinyl, 2-benzimidazolyl, 5-tetrazolyl,  
 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 1-  
 5 morpholino, 1-piperidino, or 1-pyrrolidino; and  
 $R_5$  is hydrogen or methyl.

A second group of particularly preferred compound are  
 those of the formula (I'), wherein W,  $R_1$ ,  $R_2$  and  $R_3$  are as  
 defined above, and  $R_4$  is a group of formula (D):




10

wherein  $R_8$  is methyl, ethyl, phenyl, pyridyl, isopropyl, sec-  
 butyl, tert-butyl, adamantyl, cyclopentyl, cyclohexyl,  
 cyclohexylmethyl, phenylmethyl, 3,4-(methylenedioxy)phenyl,  
 15 piperonyl, pyridylmethyl, thienylmethyl, 3-indolylmethyl, 3-  
 (N-methyl)indolylmethyl, 1,1-diphenylmethyl, naphthyl,  
 naphthylmethyl, quinolylmethyl, isoquinolylmethyl, thiazolyl,  
 thiazolylmethyl, imidazolyl, imidazolylmethyl, or a  
 derivative thereof substituted by one to three substituents  
 20 selected from chloro, fluoro, hydroxy, methoxy, and  $C_1$ - $C_4$   
 alkyl, or  $R_8$  is  $-C(CH_3)_2SCH_3$  or a sulfoxide or sulfone  
 thereof,  $-C(CH_3)_2OCH_3$ ,  $-CH_2CH_2CH_2-OCH_3$ , or  $-CH(CH_3)OH$  or a  
 tert-butyl ether thereof;  $R_9$  is either hydrogen or a group  
 selected from methyl, ethyl, phenyl, 3,4-  
 25 (methylenedioxy)phenyl, piperonyl, pyridyl, benzimidazolyl  
 and 4-tetrazolyl, which group is unsubstituted or substituted  
 by one to three substituents selected from chloro, fluoro,

17

hydroxy, methoxy, methoxy-carbonyl, ethoxycarbonyl and  $C_1$ - $C_4$  alkyl; or  $R_8$  and  $R_9$ , taken together with the nitrogen atom to which they are attached, constitute a dihydrocarbostyryl ring (D') :



5

wherein the nitrogen atom may be substituted by methyl, ethyl, propyl, methoxycarbonyl or ethoxycarbonyl, and the phenyl ring may be substituted by one or two substituents selected from chloro, fluoro, methyl, methoxy or 3,4-methylenedioxy; and  $R_5$  is hydrogen or methyl.

A third group of particularly preferred compound are those of the formula (I'), wherein W,  $R_1$ ,  $R_2$  and  $R_3$  are as defined above, and  $R_4$  and  $R_5$ , taken together with the nitrogen atom to which they are attached, constitute an azaheterocyclic ring selected from morpholine, thiomorpholine, pyrrolidine, piperidine, piperazine, pyridazine, thiazolidine, tetrahydroisoquinoline, hexamethyleneimine and hexahydropyridazine, either unsubstituted or substituted by one or more substituents selected from methyl, ethyl, phenyl, 4-fluorophenyl, benzyl, alpha-methylbenzyl, hydroxy, hydroxymethyl, and carbamoyl, or by a group -CONH-A wherein A is selected from methyl, isopropyl, tert-butyl, cyclopropyl cyclobutyl, cyclopentyl, cyclohexyl, phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 4-piperidyl,

2-thiazolyl, 1-naphthyl, 2-naphthyl, 3-quinolyl, 5-quinolyl, 3-isoquinolyl, 5-isoquinolyl, 3-quinuclidinyl, 3,4-(methylenedioxy)phenyl and piperonyl.

When  $R_1$  is a hydroxymethyl derivative, it is preferably  
 5 -CH<sub>2</sub>OCH<sub>3</sub>, -CH<sub>2</sub>-O-(tetrahydropyranyl), -CH<sub>2</sub>OCOCH<sub>3</sub>, -CH<sub>2</sub>OCOC(CH<sub>3</sub>)<sub>3</sub>, -CH<sub>2</sub>OCONH<sub>2</sub>, -CH<sub>2</sub>OCONHCH<sub>3</sub>, -CH<sub>2</sub>OCON(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>OCONH-cyclohexyl, -CH<sub>2</sub>OCON(CH<sub>3</sub>)-cyclohexyl, -CH<sub>2</sub>-O-CO-morpholino, -CH<sub>2</sub>-O-CO-pyrrolidino, -CH<sub>2</sub>-O-CO-piperidino; or  
 it is -CH<sub>2</sub>OCH<sub>2</sub>Ph, -CH<sub>2</sub>OCOPh, -CH<sub>2</sub>OCOCH<sub>2</sub>Ph, -CH<sub>2</sub>OCONHCH<sub>2</sub>Ph or a  
 10 derivative thereof wherein the Ph group is substituted by one, two or three substituents selected from chloro, fluoro, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, methyl and carbamoyl, or by 3,4-dioxymethylene.

When  $R_1$  is a mercaptomethyl derivative, it is preferably  
 15 a group -CH<sub>2</sub>S-COCH<sub>3</sub>, -CH<sub>2</sub>S-COPh, -CH<sub>2</sub>S-CH<sub>3</sub>, -CH<sub>2</sub>S-Ph, -CH<sub>2</sub>S-CH<sub>2</sub>Ph, -CH<sub>2</sub>SO<sub>2</sub>-CH<sub>3</sub>, -CH<sub>2</sub>SO<sub>2</sub>-Ph, -CH<sub>2</sub>SO<sub>2</sub>-CH<sub>2</sub>Ph, -CH<sub>2</sub>S-thienyl, -CH<sub>2</sub>S-(1-methyl-1,2,3,4-tetrazol-5-yl), or a derivative thereof wherein the Ph group is substituted by one or two  
 substituents selected from chloro, fluoro, hydroxy, C<sub>1</sub>-C<sub>4</sub>  
 20 alkoxy, methyl and carbamoyl, or by 3,4-dioxymethylene.

Preferred compounds are also those of the formula (I'), wherein


W is either -NHOH or OH;  $R_1$  is either hydroxymethyl or  
 25 mercaptomethyl, or it is a hydroxymethyl derivative as defined above, or it is a mercaptomethyl derivative as defined above;

$R_2$  is hydroxy;  $R_3$  is isobutyl, cyclopentylmethyl, 3-(4-chlorophenyl)propyl, 3-(4-methoxyphenyl)propyl, 3-(4-biphenyl)propyl, 3-[4-(4-fluorophenyl)phenyl]propyl, 3-[4-(4-chlorophenyl)phenyl]propyl, 3-[4-(4-

methoxyphenyl)phenyl]propyl, 3-(4-phenoxyphenyl)propyl, 3-(4-pyridoxyphenyl)propyl, (4-methoxybenzene)sulfonylmethyl, and (4-butoxybenzene)sulfonylmethyl; and  $R_4$  and  $R_5$  are as defined above.

5       Compounds of the general formula (I) and their salts may be prepared by any suitable method known in the art, or by the following process which forms another aspect of the present invention. In the description and formulae below, the groups  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are as defined above. It is  
10 understood that in the processes below any functional group (e.g. carboxyl, hydroxyl, mercapto or amino), if needed or desired, can be masked by conventional methods and unmasked at the end or when convenient. Suitable protecting groups for such functionalities will be apparent to those skilled in the  
15 art and are well described in the chemical literature (see, for example: "Protective Groups in Organic Synthesis" by T.W. Greene, Wiley Interscience). It is also understood that the groups  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  may be converted by conventional methods into different groups included within those  
20 previously defined, if desired, at the end or at any stage of the process below. These conversions are known or will be apparent to those skilled in the art and are well described in the chemical literature (see, for example: "Comprehensive Organic Transformation" by R.C. Larock, VCH Publishers).  
25 Accordingly, the present invention provides a process for producing a compound of the invention as defined above, which process comprises:

(a) coupling an acid of formula (II), or a salt thereof,




wherein  $R_1$ ,  $R_2$  and  $R_3$  are as defined in claim 1 and  $W'$  is protected -OH or protected -NHOH, with an amine of formula (III) or a salt thereof




5

wherein  $R_4$  and  $R_5$  are as defined above, to obtain a compound of formula (Ia)



10 wherein  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$  and  $W'$  are as defined above, and then removing the protecting group from this compound to yield a compound of formula (I) wherein  $W$  is -OH or -NHOH or a salt thereof; or

(b) converting a compound of formula (IV)




15

wherein  $W''$  is OH or  $W'$ , and  $R_3$ ,  $R_4$ ,  $R_5$  and  $W'$  are as defined above, into a compound of formula (I) or (Ia) as defined

21

above, and then, when a compound of formula (Ia) is obtained, removing the protecting group from this compound to yield a compound of formula (I) wherein W is -OH or -NHOH, or a salt thereof; or


- 5 (c) condensing a lactone of formula (V)



wherein W', R<sub>1</sub> and R<sub>3</sub> are as defined above, with an amine of formula (III) as defined above, to obtain a compound of formula (Ia) as defined above wherein R<sub>2</sub> is hydroxy, and then removing the protecting group from this compound to yield a compound of formula (I) wherein W is -OH or -NHOH and R<sub>2</sub> is hydroxy; or

- (d) condensing a lactone of formula (VI):

15



wherein W', R<sub>2</sub> and R<sub>3</sub> are as defined above, with an amine of formula (III) as defined above, to obtain a compound of formula (Ia) above wherein R<sub>1</sub> is hydroxymethyl, and then removing the protecting group from this compound to yield a compound of formula (I) wherein W is -OH or -NHOH and R<sub>1</sub> is hydroxymethyl and, if desired,

(e) converting one compound of formula (I) or (Ia) into another compound of formula (I) or (Ia) and/or converting a compound of formula (I) into a pharmaceutically acceptable salt thereof, and/or converting a salt of a compound of formula (I) into the free compound.

When in compounds of formula (Ia), (II), (IV), (V) and (VI) above W' is protected -OH, it is preferably benzyloxy, p-nitrobenzyloxy, p-methoxybenzyloxy, tert-butoxy, benzhydryloxy, trityloxy, trimethylsilyloxy, tert-butyl-  
dimethylsilyloxy, tert-butyldiphenylsilyloxy, phenyl-  
dimethylsilyloxy, allyloxy, methoxy and ethoxy. Conversion of compounds of formula (Ia) wherein W' is protected -OH, obtained by process variants (a)-(d) above, into compounds of formula (I) wherein W is -OH or salts thereof, entails  
deprotection of said protected -OH groups. This conversion is carried out by methodologies well known in the art, as generally referred to above, for the removal of protecting groups from protected carboxylic acids. A preferred conversion of this type is hydrogenolysis, especially in the presence of a palladium catalyst, in an inert organic solvent such as ethanol or DMF or the like, especially at room temperature and under atmospheric pressure or moderate pressure, which is suitable for the conversion, e.g., of benzyl and p-nitrobenzyl esters into the parent carboxylic acids. Another preferred conversion of this type is acid hydrolysis, especially by hydrochloric acid or trifluoroacetic acid, in inert organic solvents such as dichloromethane, THF, acetonitrile and the like, optionally in admixture with water, methanol or ethanol, at temperatures ranging from -20 to +40 °C. Hydrolysis by trifluoroacetic acid, in particular, is suitable for the



conversion of tert-butyl esters and p-methoxybenzyl esters into the parent carboxylic acids, and may be carried out in the presence of scavengers of the tert-butyl cation, such as anisole, methanol, ethanol and water. A third preferred  
5 conversion of this type, suitable for the deprotection of allyl esters, is treatment with a palladium(0) compound such as tetrakis-(triphenylphosphine)-palladium(0). A fourth preferred conversion of this type, suitable for the conversion of alkyl esters, e.g. methyl, ethyl, propyl, i-  
10 propyl, n-butyl esters, into the parent carboxylic acids, is saponification in an alkaline medium, e.g. by NaOH or LiOH in a mixture of water and methanol.


When in compounds of formula (Ia), (II), (IV), (V) and (VI) above W' is protected -NHOH, it is either a group -NH-  
15 O-R<sub>10</sub>, or a group -N(R<sub>11</sub>)-O-R<sub>10</sub>, wherein R<sub>10</sub> and R<sub>11</sub> are, respectively, hydroxy and amino protecting groups known in the art. Such groups are removed, by conventional methods known in the art, from a compound of formula (Ia) obtained by process variants (a)-(d) above, to obtain the desired final  
20 compound of formula (I) wherein W is -NH-OH. Preferred R<sub>10</sub> and R<sub>11</sub> groups, which may be the same or different, are those removable by hydrogenolysis or by mild hydrolysis, including, in particular, benzyl, 4-methoxybenzyl, 4-nitrobenzyl, tert-butyl, tert-butoxycarbonyl, tetrahydropyranyl, trityl,  
25 trimethylsilyl and tert-butyldimethylsilyl. Removal of such R<sub>10</sub> and R<sub>11</sub> groups from compounds of formula (Ia) wherein W' is either -NH-O-R<sub>10</sub> or -N(R<sub>11</sub>)-O-R<sub>10</sub> to obtain compounds of formula (I) wherein W is -NH-OH is performed under conditions generally known in the art for obtaining hydroxamic acids, or  
30 salts thereof, from N-protected, O-protected, or N,O-diprotected hydroxamic acid derivatives. For example, benzyl,

4-methoxybenzyl and 4-nitrobenzyl groups may be removed, preferably, by catalytic hydrogenation, preferably in the presence of a palladium catalyst such as palladium on charcoal or palladium hydroxide; 4-methoxybenzyl, trityl, 5 tert-butyl, tert-butoxycarbonyl and tetrahydropyranyl groups may be removed, preferably, by mild acid hydrolysis; trimethylsilyl and tert-butyldimethylsilyl groups may be removed by aqueous workup or by mild acid treatment. In addition, a 4-nitrobenzyl group may be removed by dissolving 10 metal reduction, preferably by zinc or iron powder, and a tert-butyldimethylsilyl group may be removed by a fluoride reagent, preferably by tetrabutylammonium fluoride.

In process variant (a), the coupling between an acid of formula (II) or a salt thereof with an amine of formula (III) 15 or a salt thereof to give a compound of formula (Ia) as defined above is carried out in a manner which is known per se in peptide chemistry. Thus, for example, the condensation can be carried out according to the well-known acid halide, acid anhydride, activated amide, mixed anhydride, or 20 activated ester method.

In a preferred procedure, the condensation is carried out according to the acid chloride method, in particular by converting the acid of formula (II) into the corresponding acid chloride by the action of oxalyl chloride or thionyl 25 chloride. In another preferred procedure, the condensation is carried out by the activated ester method. Particularly preferred activated esters are the hydroxybenzotriazolyl, hydroxysuccinyl and pentafluorophenyl esters, which are obtained by reaction of the acid with the corresponding 30 alcohol in the presence of a dehydrating agent, for example N,N'-dicyclohexyl carbodiimide (hereinafter DCC), N,N-

dimethylaminopropyl-N'-ethyl carbodiimide hydrochloride (EDC), bis(2-oxo-3-oxazolidinyl)phosphinic chloride (BOP-Cl), or 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ). Activated esters with 1-hydroxybenzotriazole (HOBT) can be  
5 obtained and reacted in situ by use, as the condensing agent, of benzotriazolyl-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP reagent), O-benzotriazol-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), and O-benzotriazol-yl-N,N,N',N'-tetramethyluronium  
10 tetrafluoroborate (TBTU). Preferred mixed anhydrides are those obtained by reacting the acid or a salt thereof with a chlorocarbonate, such as ethyl chlorocarbonate, or with an acid halide, such as pivaloyl chloride. In a still another preferred procedure, the condensation between the acid of  
15 formula (II) or a salt thereof and the amine of formula (III) or a salt thereof is carried out directly in the presence of a known peptide coupling reagent, such as DDC, EDC, BOP-Cl or EEDQ, as mentioned before. Preferred conditions for the condensation in the presence of DCC are illustrated in  
20 example 13, in the presence of DCC and HOBT in examples 4 and 9, in the presence of DCC and EDC in examples 16, 22 and 25. The amines of formula (III) are known compounds, or can be obtained from known compounds by known procedures. The succinic acid derivatives of formula (II) are novel compounds  
25 and are therefore a further object of the present invention. They can be obtained by reacting known compounds of formula (VII) or a derivative thereof protected at the terminal carboxy group, i.e. that in allylic position to the double bond:



wherein W'' and R<sub>3</sub> are as defined above, with appropriate reagents followed by removal and/or introduction of the appropriate protecting groups and /or conversion into

5 different compounds of the formula (II).

For example, the acids of formula (II) wherein W' and R<sub>3</sub> are as defined above, and R<sub>1</sub> and R<sub>2</sub>, together with the carbon atom to which they are attached, constitute a cyclopropyl or substituted cyclopropyl ring (formula B1 above), are obtained

10 by cycloaddition of diazomethane, or a substituted derivative thereof, to a vinylene compound of formula (VII) as above defined, or a derivative thereof protected at the terminal carboxy group, followed by extrusion of nitrogen and removal of the protecting group, if present. This reaction can be

15 performed by any of the conditions known in the art for the cyclopropanation of double C-C bonds. Particularly preferred conditions are treatment of a protected derivative of the vinylene compound of formula (VII), especially a benzyl or tert-butyl ester thereof, with diazomethane, in an inert

20 organic solvent, such as dichloromethane, at temperatures ranging from 0 to +40 °C, followed by irradiation (300 nm or above) until extrusion of nitrogen is complete.

The acids of formula (II) wherein W' and R<sub>3</sub> are as defined above, R<sub>1</sub> is -CH<sub>2</sub>OH and R<sub>2</sub> is -OH, can be prepared by reacting

25 a vinylene compound of formula (VII) above, or a derivative thereof protected at the terminal carboxy group, with an oxidizing agent, known *per se* as a conventional reagent in the transformation of vinylene groups (>C=CH<sub>2</sub>) into glycols

( $>\text{COH}-\text{CH}_2\text{OH}$ ), followed by removal of protecting groups, if present.

A particularly preferred oxidizing agent for such conversion is N-methylmorpholine-N-oxide, in the presence of a catalytic  
5 amount of osmium tetroxide, in an inert organic solvent, or in a mixture of an inert organic solvent and water, which is preferably aqueous acetone, at a temperatures from  $0^\circ\text{C}$  to reflux.


The acids of formula (II), or esters thereof, wherein  $\text{W}'$  and  
10  $\text{R}_3$  are as described above,  $\text{R}_2$  is hydroxy or a protected derivative thereof, and  $\text{R}_1$  is a hydroxymethyl derivative, that is an ester, an ether, a carbonate or a carbamate, can be obtained from the corresponding compounds of formula (II) wherein  $\text{R}_1$  is hydroxymethyl, or derivatives thereof protected  
15 at the terminal carboxy group, by conditions known in the art for the conversion of primary carbinols into esters, ethers, carbonates and carbamates, followed by removal of the protecting group, if present. Specific examples are given in Preparations 4 and 5. The acids of formula (II), or esters  
20 thereof, wherein  $\text{W}'$  and  $\text{R}_3$  are as described above,  $\text{R}_2$  is hydroxy or a protected derivative thereof, and  $\text{R}_1$  is a mercaptomethyl or a mercaptomethyl derivative, that is a sulfide (i.e., a thioether), a thioester, a sulfoxide or a sulfone, can be obtained from the corresponding compounds of  
25 formula (II) wherein  $\text{R}_1$  is hydroxymethyl, by conditions known in the art for the conversion of primary carbinols into the corresponding primary mercaptanes, thioesters or sulfides. A preferred condition for the conversion of compounds of  
formula (II) wherein  $\text{R}_1$  is hydroxymethyl into corresponding  
30 thioethers, that is compounds of formula (II) wherein  $\text{R}_1$  is  $\text{CH}_2-\text{S}-\text{A}$ , being A as described above, comprises protection of

the carboxy group, followed by reaction with a mercaptane of formula A-SH under Mitsunobu conditions, that is in the presence of a phosphine (expecially, triphenylphosphine) and an azodicarboxylate (expecially, diethyl azodicarboxylate), and removal of the carboxy protecting group. Sulfone and sulfoxide derivatives are preferably obtained by oxidation of the above intermediates wherein  $R_1$  is  $-\text{CH}_2-\text{S}-\text{A}$  with an oxidizing agent, known in the art for converting sulfides into sulfoxides and sulfones. Similarly, a preferred condition for the conversion of compounds of formula (II) wherein  $R_1$  is hydroxymethyl into corresponding thioesters, that is compounds of formula (II) wherein  $R_1$  is  $-\text{CH}_2-\text{S}(\text{CO})-\text{A}$ , being A as described above, comprises protection of the carboxy group, followed by reaction with a thioacid of formula A-COSH under Mitsunobu conditions, and removal of the carboxy protecting group. A preferred condition for the conversion of compounds of formula (II) wherein  $R_1$  is hydroxymethyl into corresponding mercaptanes, that is compounds of formula (II) wherein  $R_1$  is  $-\text{CH}_2-\text{SH}$ , comprises hydrolysis of the acetic thioesters thereof, that is compounds of formula (II) wherein  $R_1$  is  $-\text{CH}_2-\text{SCOCH}_3$ , in turn prepared as described above, under general conditions known in the art.

The acids of formula (II) wherein  $W'$  and  $R_3$  are as defined above, and  $R_1$  and  $R_2$ , together with the carbon atom to which they are attached, constitute an oxyrane or substituted oxyrane ring (formula B2 above), are obtained by reacting a vinylene compound of formula (VII) above, or a derivative thereof protected at the terminal carboxy group, with a peroxide, known *per se* as a conventional reagent for the epoxidation of vinylene groups ( $>\text{C}=\text{CH}_2$ ), followed by removal


of protecting groups, if present. Alternatively, these epoxides can be obtained from compounds of formula (II) wherein  $R_1$  is hydroxymethyl and  $R_2$  is hydroxy, protected at the terminal carboxy group, by intramolecular condensation  
5 under Mitsunobu conditions, that is in the presence of a phosphine (expecially, triphenylphosphine) and an azodicarboxylate (expecially, diethyl azodicarboxylate), followed by removal of the carboxy protecting group.

The vinylene intermediates of formula (VII) are either  
10 known compounds, or can be obtained from known compounds by known methods, as evident to the one skilled in the art and as illustrated herebelow. A preferred method for obtaining intermediate compounds of formula (VII) is decarboxylative methylenation, for example by reacting malonic acids of  
15 formula (VIII), or derivatives thereof protected at the terminal carboxy group:




wherein  $W'$  and  $R_3$  are as defined above, with formaldehyde, preferably in the presence of piperidine, followed by  
20 optional removal of protecting groups, if present; this reaction is illustrated in preparation 2. The intermediates of formula (VIII) are known compounds, or can be prepared from known compounds by procedures well known per se. Particularly preferred compounds of formula (VIII), for use  
25 in the preparation of the preferred compounds of formula (I) wherein the stereochemistry of the carbon atom bearing the  $R_3$

group is the one described in formula (I'), are compounds of formula (VIII'):



wherein W' and R<sub>3</sub> are as defined above, which can be prepared  
5 from their carboxy-protected derivatives of formula (IX):



wherein W' and R<sub>3</sub> are as defined above, and W''' is a  
carboxy-protecting group, preferably benzyl, 4-methoxybenzyl,  
4-nitrobenzyl, benzhydryl, tert-butyl and methyl, by removal  
10 of said carboxy-protecting group by known methods, that is by  
acid or alkaline hydrolysis, or by hydrogenolysis.


Intermediates of formula (IX) wherein R<sub>3</sub> is isobutyl are  
known compounds; analogs wherein R<sub>3</sub>, being as described  
above, is different from isobutyl can be prepared by the same  
15 described procedures, or by methods known in the art. In  
particular, intermediate compounds of formula (IX) wherein W'  
and W''' are benzyl are prepared starting from a (D)-  
aminoacid of formula H<sub>2</sub>N-CH(R<sub>3</sub>)-COOH, by a process comprising  
deaminative bromination, protection of the carboxy group,  
20 condensation with dibenzyl malonate, and removal of the  
protecting group. Alternatively, intermediates of formula  
(IX) are obtained as the racemates, starting from racemic  
natural or non-natural aminoacids, by the same procedure



detailed above, and at the end the obtained racemates are optically resolved by crystallization with *S*-(-)-1-phenylethylamine, as illustrated in Preparation 1, to provide compounds of formula (IX) as the pure stereoisomers (that is, in the *R* configuration).

In process variant (b), the conversion of a vinylene compound of formula (IV) as described above into a compound of formula (I) or (Ia) is carried out according to the same procedures described above for the conversion of an intermediate of formula (VII) into the corresponding intermediate of formula (II); that is, the vinylene group ( $>C=CH_2$ ) is, by use of the appropriate reagent, converted either into a cyclopropane ring, an oxirane ring, a group  $>C(OH)-CH_2OH$  and ether, ester, carbonate or carbamate derivatives thereof, or a group  $>C(OH)-CH_2SH$  and sulfide, sulfoxide, sulfone and thioester derivatives thereof. Some of the preferred conditions are illustrated in the Examples. Thus, the conversion of a compound of formula (IV) into a compound of formula (I) wherein  $R_1$  and  $R_2$ , taken together with the carbon atom to which they are attached, constitute a cyclopropane ring is illustrated in Example 20. The conversion of a compound of formula (IV) into a compound of formula (I) wherein  $R_1$  and  $R_2$ , taken together with the carbon atom to which they are attached, constitute an oxirane ring is illustrated in Example 21. The conversion of a compound of formula (IV) into a compound of formula (I) wherein  $R_1$  is hydroxymethyl or protected hydroxymethyl, and  $R_2$  is hydroxy or protected hydroxy, is illustrated in examples 1 and 4. The conversion of a compound of formula (IV) into a compound of formula (I) wherein  $R_1$  is a thioether derivative of  $-CH_2-SH$  (i.e., a sulfide) and  $R_2$  is hydroxy is illustrated in Example 18.

In process variant (c), condensation of a lactone of formula (V) with an amine of formula (III) to afford a compound of formula (I) or (Ia) wherein  $R_2$  is hydroxy is carried out in an inert organic solvent, optionally in the presence of an inorganic or organic base, at a temperature from 0°C to reflux temperature, optionally in the presence of a catalytic amount of a reagent used in peptide chemistry for the formation of activated esters, preferably HOBT. The inert organic solvent is typically tetrahydrofuran, acetonitrile, DMF, DMSO, dichloromethane, ethanol or mixtures thereof, and the optional inorganic or organic base is typically potassium carbonate, cesium carbonate, triethylamine or dimethylaminopyridine. Typical conditions of process variant (c) are illustrated in Example 14. The lactones of formula (V) are novel compounds and are part of the present invention. Accordingly, the present invention further provides a compound of formula (V):




wherein  $W'$ ,  $R_1$  and  $R_3$  are as defined above, for use in the preparation of a compound of formula (I) according to process variant (c) described above. The lactones of formula (V) are prepared from compounds of formula (II) as defined above wherein  $R_2$  is hydroxy and  $W'$ ,  $R_1$  and  $R_3$  are as defined above, by intramolecular condensation (dehydration) between the hydroxy group represented by  $R_2$  and the terminal carboxy group, such condensation being carried out in an inert organic solvent such as acetonitrile or dichloromethane, in


the presence of a dehydrating agent such as dicyclohexyl carbodiimide or a water-soluble carbodiimide, for instance as described in Preparation 9, or by treating said compound of formula (II) wherein  $R_2$  is hydroxy with hydroxybenzotriazole or TBTU or with carbonyldiimidazole, or by exposing said  
5 compounds of formula (II) under Mitsunobu conditions, that is with triphenylphosphine and diethyl azodicarboxylate.

In process variant (d), condensation of a lactone of formula (VI) and an amine of formula (III) is accomplished by  
10 mixing the reagents in an inert organic solvent, optionally in the presence of an inorganic or organic base, at a temperature from 0°C to reflux temperature. The inert organic solvent is typically tetrahydrofuran, acetonitrile, DMF, DMSO, dichloromethane, ethanol or mixtures thereof. The  
15 optional inorganic or organic base is typically potassium carbonate, cesium carbonate, triethylamine or dimethylaminopyridine. Example 24, Route B, typically illustrates process variant (d).

The lactones of formula (VI) are novel compounds and are part  
20 of the present invention. Accordingly, the present invention further provides a compound of formula (VI):



wherein  $W'$ ,  $R_2$  and  $R_3$  are as defined above, for use in the preparation of compounds of formula (I) according to process  
25 variant (d) detailed above. The lactones of formula (VI) are prepared from compounds of formula (IIa), or an ester thereof:



wherein  $W'$ ,  $R_2$  and  $R_3$  are as defined above, by intramolecular condensation between the hydroxymethyl group and the terminal carboxy group, or an ester thereof. This condensation is

5 usually spontaneous at room temperature, that is the lactones of formula (VI) are directly obtained from ester derivatives of compounds of formula (IIa) under conventional conditions of ester hydrolysis, as detailed above. Alternatively, such intramolecular condensation can be aided by exposing the

10 compounds of formula (IIa) to dehydrating agents, or to Mitsunobu conditions, or under conventional conditions for converting a carboxylic acid into an "activated ester" thereof, as defined above. It is to be noted that the compounds of the formula (II) the present invention also

15 encompasses the acids of formula (IIa) and the esters thereof, in particular the tert-butyl, benzyl, p-nitrobenzyl and p-methoxybenzyl esters, which may be prepared as described above. Specific examples of the preparation of said lactones are given in Preparations 6-8.

20 Part of the process of the present invention is also the conversion of one compound of formula (I) into another compound of formula (I), for instance the conversion of a compound of formula (I) wherein  $W$  is  $-OH$  or a salt thereof, obtained by any of the process variants (a) - (d), into a

25 compound of formula (I) wherein  $W$  is  $-NHOH$  or a salt thereof. This conversion is carried out by methodologies well known in the art for the general conversion of carboxylic acids or

salts thereof into hydroxamic acids or salts thereof.

Typically, a compound of formula (I) wherein W is -OH, or a salt thereof, and wherein any reactive functional group which may be present in substituents  $R_1$ - $R_4$  is protected if

- 5 necessary or convenient, is condensed with hydroxylamine or a salt thereof, or with an O-protected hydroxylamine of formula  $R_{10}$ -O-NH<sub>2</sub> or a salt thereof, or with an N,O-diprotected hydroxylamine of formula  $R_{10}$ -O-NH- $R_{11}$  or a salt thereof, wherein  $R_{10}$  and  $R_{11}$ , as stated above, are, respectively,
- 10 hydroxy- and amino-protecting groups known in the art, to obtain a compound of formula (Ia) wherein W' is either -NH-OH, -NH-OR<sub>10</sub> or -N( $R_{11}$ )-OR<sub>10</sub>, and  $R_1$ - $R_5$  are as detailed above, and wherein any reactive functional group which may be present in substituents  $R_1$ - $R_5$  is protected if necessary or
- 15 convenient. The obtained compounds of formula (Ia) specified above are then converted into the desired compounds of formula (I) by removal of said protecting groups  $R_{10}$  and  $R_{11}$ , if present, and of any protecting group which may be present on the substituents  $R_1$ - $R_5$ . Preferred  $R_{10}$  and  $R_{11}$  groups, which
- 20 may be the same or different, are those removable by hydrogenolysis or by mild hydrolysis, including, in particular, benzyl, 4-methoxybenzyl, 4-nitrobenzyl, tert-butyl, tert-butoxycarbonyl, tetrahydropyranyl, trityl, trimethylsilyl and tert-butyldimethylsilyl. Thus, an O-
- 25 protected hydroxylamine is, typically, O-benzyl-hydroxylamine, O-(4-methoxybenzyl)hydroxylamine, O-(4-nitrobenzyl)-hydroxylamine, O-trimethylsilyl-hydroxylamine, O-(tert-butyldimethylsilyl)-hydroxylamine, O-(tert-butyl)-hydroxylamine, O-(tert-butoxycarbonyl)-hydroxylamine, O-
- 30 trityl-hydroxylamine, or O-tetrahydropyranyl-hydroxylamine. An N,O-diprotected hydroxylamine is, preferably, N,O-

bis(benzyl)-hydroxylamine, N,O-bis(4-methoxybenzyl)-hydroxylamine, N,O-bis(tert-butoxycarbonyl)-hydroxylamine, N-(tert-butoxycarbonyl)-O-(tert-butyl-dimethylsilyl)-hydroxylamine, and N-(tert-butoxycarbonyl)-O-

5 (tetrahydropyranyl)-hydroxylamine. Condensation between a compound of formula (I) wherein W is -OH or a salt thereof and hydroxylamine or a protected hydroxylamine or salts thereof can be carried out in a variety of conditions. Thus, the carboxyl group of the compounds of formula (I) wherein W

10 is -OH or a salt thereof may be converted, either *in situ* or in a distinct step, into an "activated derivative" thereof, which then reacts with the hydroxylamine, protected hydroxylamines or salts thereof which are either added or already present in the reaction mixture. "Activated

15 derivatives" of carboxylic acids of formula (I) wherein W is -OH are the acid chloride, mixed anhydrides, and esters thereof. In particular, the acid chloride is obtained by reacting the acid or a salt thereof with a reagent such as oxalyl chloride or thionyl chloride; mixed anhydrides are

20 obtained by reacting the acid or a salt thereof with a chlorocarbonate, such as ethyl chlorocarbonate, or with an acid halide, such as pivaloyl chloride; esters, which are, preferably, the methyl, ethyl, pentafluorophenyl, hydroxysuccinyl, or hydroxybenzotriazolyl esters, are

25 obtained by reaction of the acid with the corresponding alcohol in the presence of a dehydrating agent, for example DCC, EDC, BOP reagent, HBTU, TBTU, EEDQ. Alternatively, carboxylic acids of formula (I) wherein W is -OH, or salts thereof, are directly condensed with hydroxylamine, protected

30 hydroxylamines or salts thereof, in the presence of condensing agents known in the art and used for peptide

coupling, preferably BOP reagent, BOP-Cl, HBTU, TBTU, DCC and EDC. Preferably, the condensation reaction is carried out in an inert organic solvent, such as DMF, THF, acetonitrile, dichloromethane or toluene at a temperature from -20 to + 60 °C, optionally in the presence of a tertiary organic base. When protected hydroxylamines are employed, the reaction affords compounds of formula (Ia) above, wherein W' is protected -NHOH. These compounds of formula (Ia) wherein W' is protected -NHOH are then converted into compounds of formula (I) wherein W is -NHOH, or salts thereof, by removing such protecting groups, as detailed above. For example, benzyl, 4-methoxybenzyl and 4-nitrobenzyl groups may be removed, preferably, by catalytic hydrogenation, preferably in the presence of a palladium catalyst such as palladium on charcoal or palladium hydroxide; 4-methoxybenzyl, trityl, tert-butyl, tert-butoxycarbonyl and tetrahydropyranyl groups may be removed, preferably, by mild acid hydrolysis; trimethylsilyl and tert-butyldimethylsilyl groups may be removed by aqueous workup or by mild acid treatment. In addition, a 4-nitrobenzyl group may be removed by dissolving metal reduction, preferably by zinc or iron powder, and a tert-butyldimethylsilyl group may be removed by a fluoride reagent, preferably by tetrabutylammonium fluoride.

Compounds of formula (I) and intermediates of formula (Ia) or (II) wherein R<sub>2</sub> is -OH and R<sub>1</sub> is -CH<sub>2</sub>OH also serve as useful intermediates in the preparation of compounds of formula (I) wherein R<sub>2</sub> is -OH and R<sub>1</sub> is either a hydroxymethyl derivative, that is an ether, an ester, a carbonate or a carbamate thereof, or it is mercaptomethyl or a mercaptomethyl derivative, that is a thioether, a thioester, a sulfide, a sulfoxide or a sulfone thereof. This

conversion can be achieved by a method comprising (i) optional protection of reactive functional groups anywhere in the molecule; (ii) conversion of the hydroxymethyl group  $R_1$  into a new group  $R_1$  to be selected among those defined above, or a protected derivative thereof, by methodologies known *per se* in the general chemistry of primary carbinols; (iii) optional removal of protecting groups, when present. More in particular, the selected class of compounds of formula (I) wherein  $R_1$  is a hydroxymethyl derivative can be obtained by using in step (ii) above reagents and conditions known *per se* to afford ethers, esters, carbonates and carbamates from primary carbinols, for example, by methods described in "Comprehensive Organic Transformation" by R.C. Larock, VCH Publishers. Similarly, the selected class of compounds of formula (I) wherein  $R_1$  is either mercaptomethyl or a mercaptomethyl derivative can be obtained by using in step (ii) above reagents and conditions known *per se* to afford mercaptanes, thioethers, thioesters, sulfoxides and sulfones from primary carbinols, either directly or by the intermediacy of first-obtained mercaptanes or thioethers.

As mentioned above, although low molecular weight MMP inhibitors, including hydroxamates of general formula (A), are already known, there is a strong need for better and diversified molecules for their use in medicine, that is in particular in the control or prevention of degenerative joint diseases such as rheumatoid arthritis and osteoarthritis, or in the management of multiple sclerosis, or in the treatment of invasive tumors. We found that the compounds of formula (I) provided by the present invention, and in particular the compounds having the preferred stereochemistry represented in formula (I'), are characterized by high inhibitory activity



on the matrix metalloproteinases, especially collagenases, gelatinases and stromelysins. For example, the following protocol was used to assess the biochemical activity of compounds of formula (I) against MMP-1, MMP-2, and MMP-3

5 (respectively, human interstitial collagenase, gelatinase A, and stromelysin-1).

BIOCHEMICAL ASSAY (Protocol A)

The *in vitro* potency of the compounds of the present invention as competitive inhibitors of selected matrix metalloproteinases was determined as described below. Human collagenase (MMP-1) was obtained as truncated recombinant enzyme encompassing residues 101-269 and did not require activation. Human gelatinase-A (MMP-2) was obtained as pro-enzyme (72 kDa) and was activated with 1 mM 4-aminophenylmercuric acetate for 30 min at 37 °C immediately prior to use. Human stromelysin-1 (MMP-3) was obtained as a recombinant pro-enzyme isolated either from *E. coli* or from baculovirus infected Sf9 insect cells and activated either by heat (1 h, 55 °C) or by 5 mg/l trypsin (30 min, 37°C) before addition of soybean trypsin inhibitor, finally removed by centrifugation.

All enzyme assays to determine values of the enzyme-inhibitor dissociation constants were performed using the peptide substrate (7-methoxycoumarin-4-yl)Acetyl-Pro-Leu-Gly-Leu-(3-[2,4-dinitrophenyl]-L-2,3-diaminopropionyl)-Ala-Arg-NH<sub>2</sub> (Mca-Pro-Leu-Gly-Leu-Dpa-Ala-Arg-NH<sub>2</sub>) (C.G. Knight et al., FEBS Lett. 296:263-266, 1992). The enzymes cleave at the Gly-Leu bond removing the internally quenching Dpa group. The release of the highly fluorescent peptide Mca-Pro-Leu was followed fluorimetrically setting the excitation wavelength

at 326 nm (bandwidth 5 nm) and the emission at 392 nm (bandwidth 20 nm).

Continuous fluorescence assay was carried out using a Perkin Elmer LS-50 Fluorescence Spectrophotometer fitted with a thermostatted four position stirring cell changer. Inhibitors were dissolved in the assay buffer (50 mM Tris/HCl pH 7.4 containing 0.15 M NaCl, 10 mM CaCl<sub>2</sub>, 0.01 mM ZnCl<sub>2</sub> and 0.05% Brij 35) thermostatted at 37 °C. The reaction was started by the addition of substrate (2 mM), followed by the addition of the enzyme (e.g. 1.0 nM collagenase, 0.04 nM gelatinase, or 3.0 nM stromelysin) and the increase in fluorescence was monitored over 30 min. The run was repeated at different concentrations of inhibitors in the range 0.1 - 10000 nM and data at inhibitor concentration in the region of their enzyme-inhibitor dissociation constants were used for calculation.

Substrate consumption was within 5% of the initial concentration and data outside this range were discarded. The Michaelis constants ( $K_m$ ) for the enzyme-substrate complexes is 70 mM or greater (C.G. Knight et al., reference above), so that the factor  $(1 + S/K_m)$  can be approximated to unity without appreciable error.

For example, the following results were obtained for some of the compounds described in Examples below:

Compound of Example 3: MMP-1,  $K_i = 2.3$  nM; MMP-2,  $K_i = 9.2$  nM; MMP-3,  $K_i = 150$  nM.

Compound of Example 5: MMP-1,  $K_i = 2.7$  nM; MMP-2,  $K_i = 0.62$  nM; MMP-3,  $K_i = 15.0$  nM.

Compound of Example 6: MMP-1,  $K_i = 67$  nM; MMP-2,  $K_i = 23$  nM; MMP-3,  $K_i = 430$  nM.

41

Compound of Example 7: MMP-1,  $K_i = 1.5$  nM; MMP-2,  $K_i = 3.1$  nM; MMP-3,  $K_i = 32$  nM.

Compound of Example 8: MMP-1,  $K_i = 45$  nM; MMP-2,  $K_i = 40$  nM; MMP-3,  $K_i = 1100$  nM.

- 5 Compound of Example 9: MMP-1,  $K_i = 2.2$  nM; MMP-2,  $K_i = 1.7$  nM; MMP-3,  $K_i = 110$  nM.

Compound of Example 11: MMP-1,  $K_i = 7.3$  nM; MMP-2,  $K_i = 4.7$  nM; MMP-3,  $K_i = 4.9$  nM.

- 10 Compound of Example 12: MMP-1,  $K_i = 6.8$  nM; MMP-2,  $K_i = 9.5$  nM; MMP-3,  $K_i = 10.0$  nM.

Compound of Example 13: MMP-1,  $K_i = 3.4$  nM; MMP-2,  $K_i = 6.5$  nM; MMP-3,  $K_i = 71$  nM.

Compound of Example 16: MMP-1,  $K_i = 1.5$  nM; MMP-2,  $K_i = 3.2$  nM; MMP-3,  $K_i = 23$  nM.

- 15 Compound of Example 20: MMP-1,  $K_i = 25$  nM; MMP-2,  $K_i = 19$  nM; MMP-3,  $K_i = 880$  nM.

Some of the compounds of formula (I) were also shown to possess high activity at inhibiting the release of TNF from several different cell lines, under different stimulation  
20 conditions. For example, the following cell-based assay was used to assess such activity:

CELLULAR ASSAY (Protocol B)

The *in vitro* potency of the compounds of the present invention as inhibitors of the release of TNF from cells was  
25 determined as described below. THP-1 cells, cultured in RPMI 1640 supplemented with 10% FCS, were distributed into 24-well plates, 1 mL of a suspension of  $1 \times 10^6$  cells/mL in each well. Compounds to be tested, dissolved in DMSO and diluted with the culture medium (1% final DMSO concentration) were added.  
30 Plates were incubated for 30 min at 37 °C in 5% CO<sub>2</sub>, and lipopolisaccharide (LPS 0111:B4, 5 microg/mL) was added as a

stimulant. After a further 4 h incubation, cells were harvested, centrifuged (2,000 rpm, 7 min), and the supernatant was collected and frozen (-20 °C) until analysis. Analysis was run by classical ELISA methodology (monoclonal anti-TNF- $\alpha$  antibody, rabbit capture polyclonal antibody, and peroxidized anti-rabbit antibody).

For example, the following results were obtained for some of the compounds described in Examples below:

Compound of Example 2:  $IC_{50}$  = 5.2 mM

10 Compound of Example 3:  $IC_{50}$  = 8.1 mM

Compound of Example 5:  $IC_{50}$  = 1.49 mM

Compound of Example 6:  $IC_{50}$  = 65.4 mM

Compound of Example 7: 90% inhibition at 100  $\mu$ M.

Further, several compounds of formula (I), in particular  
15 when  $R_2$  is hydroxy, and when  $R_1$  is hydroxymethyl, have enhanced water solubility as compared to the vast majority of MMP inhibitors of the prior art. An aqueous solubility threshold, which may vary widely but can be roughly set in the range of 1-10 mg/mL, makes systemic administration easy, and  
20 more than often is essential to good oral bioavailability. For example, aqueous solubility (at pH 7.0, 25°C) of some of the preferred compounds were as follows:

Compound of Example 3: > 15 mg/mL

Compound of Example 7: 6 mg/mL

25 Compound of Example 8: 10 mg/mL

Compound of Example 9: > 9 mg/mL

Compound of Example 11: 3.4 mg/mL

Compound of Example 12: > 10 mg/mL

Compound of Example 13: 1.3 mg/mL

30 Compound of Example 20: 5.8 mg/mL

Compound of Example 22: 1.9 mg/mL

43

Compound of Example 23: 5 mg/mL

Compound of Example 24: > 6 mg/mL

Compound of Example 25: > 8 mg/mL

The enhanced metabolic stability of the compounds of the present invention, observed in vitro toward hepatocyte suspensions, is reflected in superior pharmacokinetic properties, as determined by observation of plasma levels achievable after i.v. and oral administration to rats of compounds of formula (I) wherein W is -NHOH (Protocol C).

10 PHARMACOKINETIC ASSAY (Protocol C)

Male rats (Sprague-Dawley; body weight about 250-300 g) were acclimatized for about one week before the start of the study. Approximately one week before dosing and while under anesthesia, rats were fitted with a cannula implanted in the superior vena cava via the jugular vein. Before treatment, rats were overnight fasted. For each compound to be assessed, three rats were treated intravenously (injection in the tail vein as a bolus) with a single dose of 15 mg/kg (as free base or free acid if needed) body weight. Other three rats were treated orally (single bolus dose by gastric gavage) with 15 mg/kg of the same compound. In some cases, according to the detection limit of each particular compound, and its availability in stock, these standard doses were modified as needed.

25 Blood samples were collected into syringes especially prepared for use with the cannulated animals via the SCV cannula. Under standard conditions, blood samples (0.2 mL) were transferred into heparinized tubes at the following times: 0 (pre-dose), 2, 4, 8, 16, 30, 45, 60, 90 min, 2, 4, 6, 8 and 24 h after iv administration; 0 (pre-dose), 5, 10, 30, 45, 60, 90 min, 2, 4, 6, 8, and 24 h after oral administration. Plasma was

obtained immediately by centrifugation (about 10,000 rpm for 3 min) and stored at -80 °C till analysis. Plasma samples were assayed for parent drug concentrations using an appropriate analytical method. Under standard conditions, liquid chromatography - tandem mass spectrometry (LC/MS-MS) was used.

Non-compartmental pharmacokinetic calculations were performed with the aid of Pharm-NCA package (SIMED, France). Maximum plasma concentration,  $C_{max}$ , and the corresponding time,  $t_{max}$ , were read as the coordinates of the highest raw-data point for individual plasma profiles. Terminal phase was evaluated by graphic judgement by eye; terminal rate constant,  $l_z$ , was calculated as the slope of the terminal linear portion of natural log-transformed plasma concentrations vs. time curve, obtained using linear regression. Terminal half-life,  $t_{1/2,z}$ , was calculated as  $\ln(2)/l_z$ . The area under the plasma concentration-time curve was estimated by the linear trapezoidal rule up to the last measured concentration ( $AUC(0-t_z)$ ) and extrapolated to infinity assuming monoexponential decay, using the formula  $AUC = AUC(0-t_z) + C(t_z)/l_z$ , where  $C(t_z)$  is the measured concentration of the last point with detectable concentration. Following iv dosing, systemic clearance was obtained using the formula  $Cl = \text{Dose}/AUC$ . According to Protocol C, the pharmacokinetic in rats of some representative compounds within the present invention was evaluated. Significant pharmacokinetic data obtained in rats at a 15 mg/kg dose were as follows:

Compound of Example 3: Clearance, 9.89 mL/min/kg; AUC iv, 1.53 mg x min/mL;  $t_{1/2,z}$ , 5.4 hr; AUC os, 66 µg x min/mL.

Compound of Example 16: Clearance, 11 mL/min/kg; AUC iv, 1.35 mg x min/mL;  $t_{1/2,z}$ , 2.3 hr; AUC os, 80 µg x min/mL.

In particular, the plasma clearance of the compounds of Example 3 and Example 16 is inferior to that reported for any hydroxamic acid MMP inhibitor disclosed in the literature, to our best knowledge.

5       The novel compounds of formula (I) can be used in a method of treatment of the human or animal body by therapy. The condition of a human or animal patient may thus be improved. In particular, compounds of the present invention can be used to limit the local growth of established tumors, and  
10   to limit growth of secondary metastases. The following protocol was used to evaluate the antitumoral efficacy in vivo of the compounds of the present invention:

ANTITUMOR EFFICACY ASSAY (Protocol D):

Human DU145 Prostatic Carcinoma cells ( $2 \times 10^6$ ) were injected  
15   subcutaneously at day 0 in Balb nu/nu mice. Treatment with inhibitor (i.p. injection) started at day +1 and terminated at day +20. Animals were sacrificed at day +21, and the tumors excised and weighted. Percent tumor growth inhibition vs. control mice (%TGI) was calculated as follows:

$$20 \quad \%TGI = 100 - \frac{(\text{Mean Tumor Weight in treated group})}{(\text{Mean Tumor Weight in control group})} \times 100$$

For example, the TGI determined after 100 mg/kg b.i.d. treatment with the compound of Example 16, was 83% ( $p < 0.01$ , Dunnett method).

25       The compounds of the present invention are typically administered in the form of pharmaceutical preparations which contain them in association with a compatible pharmaceutical carrier material. Thus, a distinct aspect of the present invention is the preparation of pharmaceutical compositions  
30   carrying a compound of formula (I) as active ingredient, and a method of management (i.e. treatment or prophylaxis) of

diseases or conditions mediated in humans and warm blood animals by MMPs, which method comprises administering to the mammal an effective amount of a compound of formula (I) above, or a pharmaceutically acceptable salt thereof.

5       The present invention therefore further provides a compound of the invention as defined above for use in a method of treatment or prophylaxis of a disease mediated in a mammal by a matrix metalloproteinase, and in a method of treatment or prophylaxis of an inflammatory, infectious or  
10 immunological disease prompted by the release of TNF.

In particular, the compounds of formula (I) can be administered:

A) Orally, for example, as tablets, coated tablets, dragees, troches, lozenges, aqueous or oily suspensions,  
15 dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from  
20 the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for  
25 the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, maize starch, or alginic acid; binding agents, for example starch,  
30 gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be



uncoated or they may be coated by known techniques to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is present as such, or mixed with water or an oil medium, for example, peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinylpyrrolidone gum tragacanth and gum acacia; dispersing or wetting agents may be naturally-occurring phosphatides, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyoxyethylene sorbitan monooleate. The said aqueous suspensions may also contain one or more preservatives, for example, ethyl or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more

flavoring agents, or one or more sweetening agents, such as sucrose or saccharin.

Oily suspension may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oils, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan mono-oleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening

and flavoring agents. Syrups and elixirs may be formulated with sweetening agents, for example glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents;

5           B) Parenterally, either subcutaneously, or intravenously, or intramuscularly, or intrasternally, or by infusion techniques, in the form of sterile injectable aqueous or olagenous suspensions. This suspension may be formulated according to the known art using those suitable dispersing of  
10 wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and  
15 solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition fatty  
20 acids such as oleic acid find use in the preparation of injectables;

C) By inhalation, in the form of aerosols or solutions for nebulizers;

D) Rectally, in the form of suppositories prepared by  
25 mixing the drug with a suitable non-irratating excipient which is solid at ordinary temperature but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and poly-ethylene glycols;

30           E) Topically, in the form of creams, ointments, jellies, collyriums, solutions or suspensions.


Daily dosages can vary within wide limits and will be adjusted to the individual requirements in each particular case. In general, for administration to adults, an appropriate daily dosage is in the range of about 5 mg to about 500 mg, although the upper limit may be exceeded if expedient. The daily dosage can be administered as a single dosage or in divided dosages.

The following Examples are meant to illustrate the present invention in more detail.

10

**EXAMPLE 1**

**[(2R,3S)-3,4-Dihydroxy-2-isobutyl-3-(tetrahydropyran-2-yloxy)methyl]succinyl]-L-tert-leucine-N-methylamide**



**Step 1: [(2R)-4-Benzyloxy-3-benzyloxycarbonyl-2-isobutyl-succinyl]-L-tert-leucine-N-methylamide**

Benzyloxy (3R)-(2-benzyloxycarbonyl-3-carboxy-5-methyl)hexanoate (6.0 g, 15 mmol), obtained as described in Preparation 1, was dissolved in DMF (12 mL). To this solution, L-tert-leucine-N-methylamide (2.15 g, 15 mmol) and HOBt (1-hydroxybenzotriazole hydrate; 2.03 g, 15 mmol) were added at room temperature, followed by dropwise addition at 0°C of a solution of DCC (dicyclohexylcarbodiimide; 7.85 g, 15 mmol) in THF (12 mL). The mixture was kept at 0°C for 30 min, then kept under stirring for 5 h at room temperature. Dicyclohexylurea was filtered off, the solvent was removed in vacuo, the residue

51

was taken up in ethyl acetate, and the solution was sequentially washed with 10% aqueous citric acid (100 mL), saturated aqueous sodium bicarbonate (100 mL), and brine (100 mL). After drying over Na<sub>2</sub>SO<sub>4</sub>, the solvent was removed to give an oil. Purification by flash chromatography over silica (n-hexane and ethyl acetate gradient elution) afforded the title compound as a white foam (6.6 g, 13 mmol; 86.6%). <sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>): 0.66 and 0.71 (each d, J= 6.4 Hz, 6 H, 2 x Me), 0.82 (s, 9 H, t-Bu), 0.9-1.4 (m, 3 H, CH<sub>2</sub>CHMe<sub>2</sub>), 2.49 (d, J= 4.6 Hz, 3 H, CONHCH<sub>3</sub>), 3.15 (m, 1 H, CH-iBu), 3.65 (d, J= 9.9 Hz, 1 H, CH(CO<sub>2</sub>Bn)<sub>2</sub>), 4.09 (d, J= 9.4 Hz, 1 H, CONHCH), 5.02 (m, 4 H, 2 x CO<sub>2</sub>CH<sub>2</sub>Ph), 7.28 (m, 10 H, 2 x Ph), 7.76 (q, J= 4.6 Hz, 1 H, CONHCH<sub>3</sub>), and 7.82 ppm (d, J= 9.4 Hz, 1 H, CONHCH).

**Step 2: [(2R)-4-Hydroxy-2-isobutyl-3-methylenesuccinyl]-L-tert-leucine-N-methylamide**

A solution of the compound from Step 1 above (6.6 g, 13 mmol) in ethanol (100 mL) was treated with 10% Pd on charcoal (0.6 g) and exposed to a hydrogen atmosphere for 3 h. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol (50 mL), and the resulting solution of crude (2R)-(4-hydroxy-3-hydroxycarbonyl-2-isobutylsuccinyl)-L-tert-leucine-N-methylamide was treated with piperidine (1.5 mL, 15 mmol) and 37% aqueous formaldehyde (10.1 mL, 0.136 mol). After 20 h at room temperature, the mixture was evaporated under reduced pressure and the residue partitioned between ethyl acetate and 4% aqueous HCl. The organic layer was sequentially washed with brine and water, then dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated, to afford the title compound as a white solid (3.16 g, 10 mmol; 80.3%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.75 and 0.80 (each d, J= 6.4 Hz, 6 H, 2 x Me), 0.78 (s, 9 H,

52

t-Bu), 1.3-1.6 (m, 3 H,  $\text{CH}_2\text{CHMe}_2$ ), 2.51 (d,  $J = 4.7$  Hz, 3 H,  $\text{CONHCH}_3$ ), 3.54 (m, 1 H,  $\text{CH-iBu}$ ), 4.10 (d,  $J = 9.4$  Hz, 1 H,  $\text{CONHCH}$ ), 5.65 and 6.10 (each s, 2 H,  $\text{C=CH}_2$ ), 7.36 (d,  $J = 9.4$  Hz, 1 H,  $\text{CONHCH}$ ), 7.87 (q,  $J = 9.4$  Hz, 1 H,  $\text{CONHCH}_3$ ), and 12.7  
5 ppm (broad s, 1 H,  $\text{CO}_2\text{H}$ ).

**Step 3: [(2R,3S)-4-Benzhydryloxy-3-hydroxy-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-methylamide**

A solution of the compound from Step 2 above (3.16 g, 10 mmol) in tert-butanol (35 mL) was added to a mixture of N-methylmorpholine-N-oxide (60% in water; 3.32 g, 17 mmol),  
10 osmium tetroxide (2.5% in tert-butanol; 0.813 mL, 0.08 mmol), acetone (1 mL) and water (2 mL). After stirring for 20 h at room temperature, the solution was evaporated to dryness to leave crude (3S,2R)-(3,4-dihydroxy-3-hydroxymethyl-2-isobutylsuccinyl)-L-tert-leucine-N-methylamide as a light tan foam.  
15 This material was dissolved in acetonitrile, and diphenyl diazomethane (DDM; 1.95 g, 10 mmol) was added portionwise. After 2 h at room temperature, glacial acetic acid was added dropwise until the colour of excess DDM was discharged.  
20 Removal of the solvent and flash chromatography over silica (n-hexane and ethyl acetate gradient elution) afforded the title compound as the major pure diastereoisomer (2.7 g, 5.3 mmol; 52.7%) as a white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO-d}_6$ ): 0.54 and 0.61 (each d,  $J = 6.4$  Hz, 6 H, 2 x Me), 0.60 and 1.61 (each  
25 m, 2 H,  $\text{CH}_2\text{CHMe}_2$ ), 0.88 (s, 9 H, t-Bu), 1.22 (m, 1 H,  $\text{CHMe}_2$ ), 2.54 (d,  $J = 4.7$  Hz, 3 H,  $\text{CONHCH}_3$ ), 2.88 (dd,  $J = 3.5$  and 12.0 Hz, 1 H,  $\text{CH-iBu}$ ), 3.42 and 3.78 (each m, 2 H,  $\text{CH}_2\text{OH}$ ), 4.13 (d,  $J = 9.1$  Hz, 1 H,  $\text{CONHCH}$ ), 4.93 (dd,  $J = 4.4$  and 7.3 Hz, 1 H,  $\text{CH}_2\text{OH}$ ), 5.41 (s, 1 H, tertiary OH), 6.83 (s, 1 H,  $\text{CHPh}_2$ ), 7.2-  
30 7.4 (m, 10 H, 2 x Ph), 7.86 (d,  $J = 9.1$  Hz, 1 H,  $\text{CONHCH}$ ), and 7.94 ppm (q,  $J = 4.7$  Hz, 1 H,  $\text{CONHCH}_3$ ).

**Step 4: [(2R,3S)-4-Benzhydryloxy-3-hydroxy-2-isobutyl-3-(tetrahydropyran-2-yloxymethyl)succinyl]-L-tert-leucine-N-methylamide**

A solution of the compound from Step 3 above (2.7 g, 5.3 mmol) in dichloromethane (50 mL) was treated at 0°C with pyridinium-p-toluenesulfonate (150 mg) and 3,4-dihydro-2H-pyran (4.8 mL, 5.3 mmol). The mixture was kept in a cold room (4°C) for 3 days, then diluted with ethyl acetate (150 mL) and sequentially washed with 4% aqueous HCl, saturated aqueous sodium bicarbonate, and water. After drying over Na<sub>2</sub>SO<sub>4</sub> and evaporation, a colourless oil was obtained, which was treated with petroleum ether (150 mL) and kept at 4°C overnight. A white precipitate formed, which was recovered by filtration, thereby obtaining the title compound (2.1 g, 3.5 mmol; 66%) as a diastereoisomeric mixture at the 2-tetrahydropyran carbon atom. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.59 and 0.65 (each d, J= 6.7 Hz, 6 H, 2 x Me), 0.7-1.7 (m, 9 H, CH<sub>2</sub>CHMe<sub>2</sub> and pyran 3,4,5-methylene), 0.87 (s, 9 H, t-Bu), 2.53 (d, J= 4.4 Hz, 3 H, CONHCH<sub>3</sub>), 2.90 (m, 1 H, CH-iBu), 3.1-3.7 (m, 2 H, pyran 6-methylene), 3.30 and 3.98 (each d, J= 10.2 Hz, CH<sub>2</sub>O-pyran of pyran isomer A), 3.60 and 3.98 (each d, J= 10.0 Hz, CH<sub>2</sub>O-pyran of pyran isomer B), 4.13 and 4.15 (each d, J= 9.4 Hz, 1 H, CONHCH of the two isomers at pyran), 4.26 and 4.47 (each m, 1 H, anomeric CH of two pyran isomers), 5.63 and 5.72 (each s, 1 H, tertiary OH of the two isomers at pyran), 6.87 (s, 1 H, CHPh<sub>2</sub>), 7.2-7.4 (m, 10 H, 2 x Ph), 7.84 and 7.85 (each d, J= 9.4 Hz, 1 H, CONHCH of the two isomers at pyran), and 7.95 ppm (q, J= 4.4 Hz, 1 H, CONHCH<sub>3</sub>).

**Step 5: [(2R,3S)-3,4-Dihydroxy-2-isobutyl-3-(tetrahydropyran-2-yloxymethyl)-succinyl]-L-tert-leucine-N-methylamide**

54


The compound from Step 4 above (2.1 g, 3.5 mmol) was taken up in ethanol (50 mL) and exposed to a hydrogen atmosphere in the presence of 10% Pd on charcoal (0.2 g) overnight. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol, and the combined ethanol solution was evaporated to dryness. The residue was taken up with n-hexane, stirred for 30 min, and the resulting white powder was collected by filtration, thereby obtaining the title compound (diastereomeric mixture at the anomeric tetrahydropyran carbon atom; 1.39 g, 3.2 mmol; 92%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.80 (d, J= 6.5 Hz, 6 H, 2 x Me), 0.87 (s, 9 H, t-Bu), 0.9-1.7 (m, 9 H, CH<sub>2</sub>CHMe<sub>2</sub> and pyran 3,4,5-methylene), 2.53 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>), 2.71 (m, 1 H, CH-iBu), 3.21 and 3.80 (each d, J= 10 Hz, CH<sub>2</sub>O-pyran of pyran isomer A), 3.52 (m, CH<sub>2</sub>O-pyran of pyran isomer B), 3.30 and 3.70 (each m, 2 H, pyran 6-methylene), 4.10 and 4.11 (each d, J= 9.4 Hz, 1 H, CONHCH of the two isomers at pyran), 4.45 and 4.49 (each m, 1 H, anomeric CH of two pyran isomers), 5.40 (broad s, 1 H, tertiary OH), 7.66 and 7.67 (each d, J= 9.4 Hz, 1 H, CONHCH of the two isomers at pyran), and 7.90 ppm (q, J= 4.7 Hz, 1 H, CONHCH<sub>3</sub>).

**EXAMPLE 2**

**[(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(tetrahydropyran-2-yloxymethyl)succinyl]-L-tert-leucine-N-methylamide**



55



**Step 1: [(2R,3S)-4-Benzyloxyamino-3-hydroxy-2-isobutyl-3-(tetrahydropyran-2-yloxymethyl)succinyl]-L-tert-leucine-N-methylamide**

- 5 The compound described in Example 1 above (1.39 g, 3.2 mmol) was dissolved in acetonitrile (50 mL). To this solution, O-benzylhydroxylamine hydrochloride (0.59 g, 3.63 mmol), N-methylmorpholine (0.8 mL, 7.35 mmol), and O-benzotriazol-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU; 1.29 g, 4.01 mmol) were added in this order. After stirring for 20 h at room temperature, the solvent was removed under reduced pressure and the residue was taken up in ethyl acetate. Sequential washing with 2N aqueous HCl, saturated aqueous sodium bicarbonate and water, followed by drying over Na<sub>2</sub>SO<sub>4</sub> and evaporation, left a residue which was purified by flash chromatography over silica (n-hexane and ethyl acetate gradient elution), thereby affording the two isomers (at the pyran anomeric C) of the title product as white solids (sum of the two 740 mg, 1.38 mmol; 42%). Isomer A, <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.73 and 0.76 (each d, J= 6.7 Hz, 6 H, 2 x Me), 0.89 (s, 9 H, t-Bu), 0.9-1.6 (m, 9 H, CH<sub>2</sub>CHMe<sub>2</sub> and pyran 3,4,5-methylene), 2.53 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>), 2.89 (m, 1 H, CH-iBu), 3.11 and 3.91 (each d, J= 10.3 Hz, 2H, CH<sub>2</sub>O-pyran),
- 10
- 15
- 20

56

3.30 and 3.65 (each m, 2 H, pyran 6-methylene), 4.12 (d, J= 9.1 Hz, 1 H, CONHCH), 4.52 (m, 1 H, anomeric CH of pyran), 4.78 (m, 2 H, Ph-CH<sub>2</sub>O), 5.33 (s, 1 H, tertiary OH), 7.35 (m, 5 H, Ph), 7.84 (q, J= 4.7 Hz, 1 H, CONHCH<sub>3</sub>), 8.06 (d, J= 9.1 Hz, 1 H, CONHCH), and 11.20 ppm (s, 1 H, CONHO). Isomer B, <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 1.17 and 1.20 (each d, J= 6.7 Hz, 6 H, 2 x Me), 1.32 (s, 9 H, t-Bu), 1.1-2.1 (m, 9 H, CH<sub>2</sub>CHMe<sub>2</sub> and pyran 3,4,5-methylene), 2.97 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>), 3.27 (dd, J= 3.5 and 11.7 Hz, 1 H, CH-iBu), 3.80 and 4.10 (each m, 2 H, pyran 6-methylene), 3.87 and 4.12 (each d, J= 10.2 Hz, 2H, CH<sub>2</sub>O-pyran), 4.57 (d, J= 9.1 Hz, 1 H, CONHCH), 4.87 (m, 1 H, anomeric CH of pyran), 5.20 (s, 2 H, Ph-CH<sub>2</sub>O), 5.79 (s, 1 H, tertiary OH), 7.78 (m, 5 H, Ph), 8.31 (q, J= 4.7 Hz, 1 H, CONHCH<sub>3</sub>), 8.46 (d, J= 9.1 Hz, 1 H, CONHCH), and 11.43 ppm (s, 1 H, CONHO).

**Step 2: [(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(tetrahydropyran-2-yloxymethyl)succinyl]-L-tert-leucine-N-methylamide**

A solution of the compound from Step 1 above (combined mixture of isomers A and B; 740 mg, 1.38 mmol) in ethanol (40 mL) was exposed to a hydrogen atmosphere in the presence of 10% Pd on charcoal (74 mg) for 2 h. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol, and the combined ethanol solution was evaporated to dryness. The residue was taken up with n-hexane, stirred for 30 min, and the resulting white powder was collected by filtration, thereby obtaining the title compound (diastereomeric mixture at the 2-tetrahydropyran carbon atom; 565 mg, 1.27 mmol; 92%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.78 (m, 6 H, 2 x Me), 0.89 (s, 9 H, t-Bu), 0.9-1.8 (m, 9 H, CH<sub>2</sub>CHMe<sub>2</sub> and pyran 3,4,5-methylene), 2.54 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>),


57

2.80 and 2.86 (each dd,  $J = 3.5$  and  $11.7$  Hz, 1 H,  $\text{CH-iBu}$  of two pyran isomers), 3.12 and 3.89 (each d,  $J = 10.2$  Hz,  $\text{CH}_2\text{O}$ -pyran of pyran isomer A), 3.41 and 3.68 (each d,  $J = 10.2$  Hz,  $\text{CH}_2\text{O}$ -pyran of pyran isomer B), 3.30-3.60 (two m, 2 H, pyran 6-methylene), 4.12 (d,  $J = 9.1$  Hz, 1 H,  $\text{CONHCH}$ ), 4.41 and 4.50 (each m, 1 H, anomeric CH of two pyran isomers), 5.23 and 5.29 (each s, 1 H, tertiary OH of two pyran isomers), 7.85 and 7.88 (each q,  $J = 4.7$  Hz, 1 H,  $\text{CONHCH}_3$  of two pyran isomers), 7.94 and 7.99 (each d,  $J = 9.1$  Hz, 1 H,  $\text{CONHCH}$  of two pyran isomers), 8.72 and 8.73 (each s, 1 H,  $\text{NHOH}$  of two pyran isomers), and 10.27 and 10.39 ppm (each s, 1 H,  $\text{NHOH}$  of two pyran isomers).

By the same procedure, applied to each separate isomer (A and B) of the compound of Step 1 above, the separate isomers (A and B) of the title compound were obtained.

### EXAMPLE 3

**[(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutyl]-succinyl]-L-tert-leucine-N-methylamide**



20


The compound described in Example 2 above (570 mg, 1.28 mmol) was dissolved in 80% aqueous ethanol (40 mL), and pyridinium p-toluenesulfonate (306 mg, 1.22 mmol) was added. The solution was heated to  $50^{\circ}\text{C}$  for 3 h, the solvent was removed under reduced pressure and the residue was purified by reverse phase chromatography over silica (water and ethanol gradient

elution) to afford the title product as a white solid (372 mg, 1.03 mmol; 85%).  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO-d}_6$ ): 0.76 and 0.78 (each d,  $J = 6.4$  Hz, 6 H, 2 x Me), 0.89 (s, 9 H, t-Bu), 0.97 and 1.62 (each m, 2 H,  $\text{CH}_2\text{CHMe}_2$ ), 1.30 (m, 1 H,  $\text{CH}_2\text{CHMe}_2$ ), 2.53 (d,  $J = 4.4$  Hz, 3 H,  $\text{CONHCH}_3$ ), 2.81 (dd,  $J = 3.2$  and 11.7 Hz, 1 H,  $\text{CH-iBu}$ ), 3.12 and 3.62 (each m, 2 H,  $\text{CH}_2\text{OH}$ ), 4.11 (d,  $J = 9.4$  Hz, 1 H,  $\text{CONHCH}$ ), 4.65 (dd,  $J = 4.7$  and 7.3 Hz, 1 H,  $\text{CH}_2\text{OH}$ ), 5.12 (s, 1 H, tertiary OH), 7.87 (m, 2 H,  $\text{CONHCH}_3$  and  $\text{CONHCH}$ ), 8.70 (s, 1 H,  $\text{NHOH}$ ), and 10.21 ppm (s, 1 H,  $\text{NHOH}$ ).

10

**EXAMPLE 4**

***[(2R,3S)-3,4-Dihydroxy-2-isobutyl-3-(tetrahydropyran-2-yloxyethyl)-succinyl]-L-phenylalanine-N-methylamide***



***Step 1: [(2R)-4-Benzyloxy-3-benzyloxycarbonyl-2-isobutyl-succinyl]-L-phenylalanine-N-methylamide***

Benzyl (3R)-(2-benzyloxycarbonyl-3-carboxy-5-methyl)hexanoate (33.5 g, 84 mmol), obtained as described in Preparation 1, was dissolved in DMF (100 mL). To this solution, L-phenylalanine-N-methylamide hydrochloride (21.1 g, 84 mmol), HOBT (1-hydroxybenzotriazole hydrate; 11 g, 81 mmol) and triethylamine (11.7 mL, 84 mmol) were added at room temperature, followed by dropwise addition at 0°C of a solution of DCC (dicyclohexylcarbodiimide; 17.33 g, 84 mmol) in THF (100 mL).

The mixture was kept at 0°C for 30 min, then kept under stirring for 5 h at room temperature. Dicyclohexylurea was filtered off, the solvent was removed in vacuo, the residue was taken up in ethyl acetate, and the solution was sequentially washed with 10% aqueous citric acid (800 mL), saturated aqueous sodium bicarbonate (800 mL), and brine (800 mL). After drying over Na<sub>2</sub>SO<sub>4</sub>, the solvent was removed to give an oil, which was taken up in diisopropyl ether (200 mL). After stirring for 20 h, the title compound was collected by filtration as a white solid (38.0 g, 68 mmol; 81%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): 0.74 and 0.75 (each d, J= 6.4 Hz, 6 H, 2 x Me), 1.01 and 1.53 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.32 (m, 1 H, CHMe<sub>2</sub>), 2.65 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>), 2.92 (m, 1 H, CH-iBu), 2.99 (m, 2 H, Phe methylene), 3.79 (d, J= 9.1 Hz, 1 H, CH(CO<sub>2</sub>Bn)<sub>2</sub>), 4.48 (m, 1 H, CONHCH), 5.10 (m, 4 H, 2 x CO<sub>2</sub>CH<sub>2</sub>Ph), 5.70 (m, 1 H, CONHCH<sub>3</sub>), 6.56 (d, J= 7.6 Hz, 1 H, CONHCH), and 7.2-7.4 ppm (m, 15 H, 3 x Ph).

**Step 2: [(2R)-4-Hydroxy-2-isobutyl-3-methylenesuccinyl]-L-phenylalanine-N-methylamide**

A solution of the compound from Step 1 above (8.7 g, 14.6 mmol) in ethanol (100 mL) was treated with 10% Pd on charcoal (0.8 g) and exposed to a hydrogen atmosphere for 3 h. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol (50 mL), and the resulting solution of crude (2R)-(4-hydroxy-3-hydroxycarbonyl-2-isobutylsuccinyl)-L-phenylalanine-N-methylamide was treated with piperidine (1.7 mL, 17.5 mmol) and 37% aqueous formaldehyde (11.7 mL, 156 mmol). After 20 h at room temperature, the mixture was evaporated under reduced pressure and the residue partitioned between ethyl acetate and 4% aqueous HCl. The organic layer was sequentially washed with

brine and water, then dried over  $\text{Na}_2\text{SO}_4$  and evaporated, to leave a residue which was treated with diisopropyl ether (50 mL). Filtration afforded the title compound as a white solid (4.7 g, 12.4 mmol; 85%).  $^1\text{H}$  NMR (200 MHz,  $\text{DMSO}-d_6$ ): 0.80 (d, J= 6.7 Hz, 6 H, 2 x Me), 1.2-1.8 (m, 3 H,  $\text{CH}_2\text{CHMe}_2$ ), 2.55 (d, J= 4.5 Hz, 3 H,  $\text{CONHCH}_3$ ), 2.83 (m, 2 H, Phe methylene), 3.45 (dd, J= 5.8 and 8.4 Hz, 1 H,  $\text{CH}-i\text{Bu}$ ), 4.41 (m, 1 H,  $\text{CONHCH}$ ), 5.44 and 5.98 (each m, 2 H,  $\text{C}=\text{CH}_2$ ), 7.20 (m, 5 H, Ph), 7.76 (q, J= 4.5 Hz, 1 H,  $\text{CONHCH}_3$ ), and 7.85 ppm (d, J= 8.4 Hz, 1 H,  $\text{CONHCH}$ ).

**Step 3: [(2R,3S)-4-Benzhydryloxy-3-hydroxy-3-hydroxymethyl-2-isobutylsuccinyl]-L-phenylalanine-N-methylamide, and [(2R,3R)-4-Benzhydryloxy-3-hydroxy-3-hydroxymethyl-2-isobutylsuccinyl]-L-phenylalanine-N-methylamide**

A solution of the compound from Step 2 above (2.0 g, 5.77 mmol) in tert-butanol (20 mL) was added to a mixture of N-methylmorpholine-N-oxide (60% in water; 1.96 g, 9.8 mmol), osmium tetroxide (2.5% in tert-butanol; 0.521 mL, 0.05 mmol), acetone (1 mL) and water (2 mL). After stirring for 20 h at room temperature, the solution was evaporated to dryness to leave crude (3,4-dihydroxy-3-hydroxymethyl-2-isobutylsuccinyl)-L-phenylalanine-N-methylamide as a light tan foam. This material was dissolved in acetonitrile, and diphenyl diazomethane (DDM; 1.12 g, 5.77 mmol) was added portionwise. After 2 h at room temperature, glacial acetic acid was added dropwise until the colour of excess DDM was discharged. Removal of the solvent left a residue, which was collected as a white solid after trituration with diisopropyl ether, filtration, and washing with ethyl ether, thereby obtaining the title product, (2R,3S) isomer (1.75 g, 3 mmol; 52.8%).  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ ): 0.45 and 0.51 (each d, J= 6.7 Hz, 6 H,

61

2 x Me), 0.44 and 1.56 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 0.98 (m, 1 H, CHMe<sub>2</sub>), 2.54 (d, J= 4.4 Hz, 3 H, CONHCH<sub>3</sub>), 2.67 (dd, J= 3.2 and 12.3 Hz, 1 H, CH-iBu), 2.77 and 2.95 (each m, 2 H, Phe methylene), 3.23 and 3.64 (each m, 2 H, CH<sub>2</sub>OH), 4.46 (m, 1 H, CONHCH), 4.82 (dd, J= 4.4 and 7.8 Hz, 1 H, CH<sub>2</sub>OH), 5.32 (s, 1 H, tertiary OH), 6.81 (s, 1 H, CHPh<sub>2</sub>), 7.0-7.4 (m, 15 H, 3 x Ph), 7.87 (q, J= 4.4 Hz, 1 H, CONHCH<sub>3</sub>), and 7.96 ppm (d, J= 8.5 Hz, 1 H, CONHCH).

The mother liquors were evaporated, and the residue purified by flash chromatography over silica (n-hexane and ethyl acetate gradient elution) to afford the title product, (2R,3R) isomer (495 mg, 0.86 mmol; 14.9%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.60 and 0.64 (each d, J= 6.0 Hz, 6 H, 2 x Me), 1.0-1.5 (m, 3 H, CH<sub>2</sub>CHMe<sub>2</sub>), 2.43 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>), 2.64 (m, 1 H, CH-iBu), 2.75 and 2.92 (each m, 2 H, Phe methylene), 3.55 (m, 2 H, CH<sub>2</sub>OH), 4.36 (m, 1 H, CONHCH), 4.82 (t, J= 5.5 Hz, 1 H, CH<sub>2</sub>OH), 5.53 (s, 1 H, tertiary OH), 6.63 (s, 1 H, CHPh<sub>2</sub>), 7.0-7.4 (m, 15 H, 3 x Ph), 7.57 (q, J= 4.7 Hz, 1 H, CONHCH<sub>3</sub>), and 7.89 ppm (d, J= 8.5 Hz, 1 H, CONHCH).

**Step 4: [(2R,3S)-4-Benzhydryloxy-3-hydroxy-2-isobutyl-3-(tetrahydropyran-2-yloxymethyl)succinyl]-L-phenylalanine-N-methylamide**

A solution of the compound from Step 3 above ((2R,3S) isomer; 820 mg, 1.43 mmol) in dichloromethane (25 mL) was treated at 0°C with pyridinium-p-toluenesulfonate (50 mg) and 3,4-dihydro-2H-pyran (1.3 mL, 14.3 mmol). The reaction was complete after stirring for 20 h at room temperature. The mixture was concentrated under reduced pressure, then taken up in ethyl acetate (100 mL) and sequentially washed with 2N aqueous HCl, saturated aqueous sodium bicarbonate, and water. After drying over Na<sub>2</sub>SO<sub>4</sub> and evaporation, a solid was

obtained, which was treated with petroleum ether (50 mL) and stirred for 30 min. A white precipitate formed, which was recovered by filtration, thereby obtaining the title compound (0.7 g, 1.06 mmol; 74.3%) in the absolute (2R,3S) configuration and as a diastereomeric mixture at the anomeric tetrahydropyran carbon atom.


**Step 5: [(2R,3S)-3,4-Dihydroxy-2-isobutyl-3-(tetrahydropyran-2-yloxymethyl)-succinyl]-L-phenylalanine-N-methylamide**

The compound from Step 4 above (700 mg, 1.06 mmol) was taken up in ethanol (25 mL) and exposed to a hydrogen atmosphere in the presence of 10% Pd on charcoal (70 mg) for 2 h. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol, and the combined ethanol solution was evaporated to dryness. The residue was taken up with n-hexane, stirred for 30 min, and the resulting white powder was collected by filtration, thereby obtaining the title compound (diastereomeric mixture at the anomeric tetrahydropyran carbon atom; 470 mg, 1.0 mmol; 94.3%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.65 (m, 6 H, 2 x Me), 0.8-1.8 (m, 9 H, CH<sub>2</sub>CHMe<sub>2</sub> and pyran 3,4,5-methylene), 2.50 (m, 1 H, CH-iBu), 2.53 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>), 2.76 and 2.93 (each m, 2 H, Phe methylene), 2.98 and 3.69 (each d, J= 10.0 Hz, CH<sub>2</sub>O-pyran of pyran isomer A), 3.36 and 3.42 (each d, J= 10.0 Hz, CH<sub>2</sub>O-pyran of pyran isomer B), 3.30 and 3.60 (each m, 2 H, pyran 6-methylene), 4.34 and 4.40 (each m, 1 H, anomeric CH of two pyran isomers), 4.43 (m, 1 H, CONHCH of the two isomers at pyran), 5.83 (broad s, 1 H, tertiary OH), 7.20 (m, 5 H, Ph), 7.83 (m, 2 H, CONHCH and CONHCH<sub>3</sub>), and 12.0 ppm (broad s, 1 H, COOH).



## EXAMPLE 5

**[(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(tetrahydropyran-2-yloxymethyl)succinyl]-L-phenylalanine-N-methylamide**



**5 Step 1: [(2R,3S)-4-Benzoyloxyamino-3-hydroxy-2-isobutyl-3-tetrahydropyran-2-yloxymethyl)succinyl]-L-phenylalanine-N-methylamide**

The compound described in Example 4 above (430 mg, 0.92 mmol) was dissolved in acetonitrile (20 mL). To this solution, O-benzylhydroxylamine hydrochloride (161 mg, 1.01 mmol), N-methylmorpholine (0.222 mL, 2.02 mmol), and O-benzotriazol-yl-N,N,N',N'-tetramethyl-uronium tetrafluoroborate (TBTU; 354 mg, 1.1 mmol) were added in this order. After stirring for 20 h at room temperature, the solvent was removed under reduced pressure and the residue was taken up in ethyl acetate. Sequential washing with 2N aqueous HCl, saturated aqueous sodium bicarbonate and water, followed by drying over Na<sub>2</sub>SO<sub>4</sub> and evaporation, left a residue which was purified by flash chromatography over silica (n-hexane and ethyl acetate gradient elution), thereby obtaining the title product (diastereomeric mixture at the pyran anomeric C) as a white solid.

**Step 2: [(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(tetrahydro-pyran-2-ylloxymethyl)succinyl]-L-phenylalanine-N-methylamide**

A solution of the compound from Step 1 above (100 mg, 0.17 mmol) in ethanol (5 mL) was exposed to a hydrogen atmosphere in the presence of 10% Pd on charcoal (10 mg) for 2 h. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol, and the combined ethanol solution was evaporated to dryness. The residue was taken up with n-hexane, stirred for 30 min, and the resulting white powder was collected by filtration, thereby obtaining the title compound (diastereomeric mixture at the anomeric tetrahydropyran carbon atom; 77 mg, 0.16 mmol; 94%). For analytical purpose, the two diastereoisomers were separated by reverse phase chromatography.

ISOMER A, <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.73 and 0.74 (each d, J= 7.0 Hz, 6 H, 2 x Me), 0.90 and 1.50 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.20 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.2-1.7 (m, 6 H, pyran 3,4,5 methylene), 2.55 (d, J= 4.4 Hz, 3 H, CONHCH<sub>3</sub>), 2.60 (m, 1 H, CH-iBu), 2.80 and 2.95 (each m, 2 H, Phe methylene), 3.20 and 3.53 (each d, J= 10.6 Hz, CH<sub>2</sub>O-pyran), 3.35 and 3.72 (two m, 2 H, pyran 6-methylene), 4.34 (m, 1 H, anomeric pyran CH), 4.50 (m, 1 H, CONHCH), 5.06 (broad s, 1 H, tertiary OH), 7.1-7.3 (m, 5 H, Ph), 7.74 (broad s, 1 H, CONHCH<sub>3</sub>), 8.05 (broad s, 1 H, CONHCH), 8.60 (broad s, 1 H, CONHOH), and 10.10 ppm (broad s, 1 H, CONHOH).


ISOMER B, <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.72 and 0.74 (each d, J= 6.7 Hz, 6 H, 2 x Me), 0.80 and 1.50 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.20 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.3-1.7 (m, 6 H, pyran 3,4,5 methylene), 2.55 (d, J= 4.4 Hz, 3 H, CONHCH<sub>3</sub>), 2.60 (m, 1 H, CH-iBu), 2.80 and 2.92 (each m, 2 H, Phe methylene), 2.67 and

65

3.67 (each d,  $J = 10.0$  Hz,  $\text{CH}_2\text{O-pyran}$ ), 3.30 and 3.70 (two m, 2 H, pyran 6-methylene), 4.35 (m, 1 H, anomeric pyran CH), 4.49 (m, 1 H,  $\text{CONHCH}$ ), 5.09 (broad s, 1 H, tertiary OH), 7.1-7.3 (m, 5 H, Ph), 7.88 (q,  $J = 4.4$  Hz, 1 H,  $\text{CONHCH}_3$ ), 8.31 (d,  $J = 8.0$  Hz, 1 H,  $\text{CONHCH}$ ), 8.65 (broad s, 1 H,  $\text{CONHOH}$ ), and 10.22 ppm (broad s, 1 H,  $\text{CONHOH}$ ).

## EXAMPLE 6

**[(2R,3R)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(tetrahydropyran-2-yloxymethyl)succinyl]-L-phenylalanine-N-methylamide**



10

Starting from the (2R,3R) isomer of the compound described in Example 4 above, Step 3, and by identical procedures as those described in the Step 4, Step 5 of the same Example 4, and Step 1, Step 2 of Example 5, the title compound was obtained as a white powder.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO-d}_6$ ): 0.65 (d,  $J = 6.4$  Hz, 6 H, 2 x Me), 0.9-1.8 (m, 9 H,  $\text{CH}_2\text{CHMe}_2$  and pyran 3,4,5-methylene), 2.50 (m, 1 H,  $\text{CH-iBu}$ ), 2.51 (d,  $J = 4.7$  Hz, 3 H,  $\text{CONHCH}_3$ ), 2.80 and 2.96 (each m, 2 H, Phe methylene of two pyran isomers), 3.23 and 3.61 (each d,  $J = 10.7$  Hz,  $\text{CH}_2\text{O-pyran}$  of pyran isomer A), 3.40 and 3.55 (each d,  $J = 10.7$  Hz,  $\text{CH}_2\text{O-pyran}$  of pyran isomer B), 3.32 and 3.70 (two m, 2 H, pyran 6-methylene), 4.36 (m, 1 H,  $\text{CONHCH}$ ), 4.36 and 4.46 (each m, 1 H, anomeric CH of two pyran isomers), 5.16 and 5.19 (each


66

s, 1 H, tertiary OH of two pyran isomers), 7.0-7.2 (m, 5 H, Ph), 7.91 (broad s, 1 H, CONHCH<sub>3</sub>), 8.23 and 8.25 (each d, J= 7.5 Hz, 1 H, CONHCH of two pyran isomers), 8.81 (broad s, 1 H, NHOH), and 10.60 ppm (broad s, 1 H, NHOH).

5

**EXAMPLE 7**


**[(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutyl-succinyl]-L-phenylalanine-N-methylamide**



10 The compound described in Example 5 above (64 mg, 0.133 mmol) was dissolved in 80% aqueous ethanol (5 mL), and pyridinium p-toluenesulfonate (64 mg, 0.127 mmol) was added. The solution was heated to 50°C for 3 h, the solvent was removed under reduced pressure and the residue was purified by reverse phase  
15 chromatography over silica (water and ethanol gradient elution) to afford the title product as a white solid (45.8 mg, 0.116 mmol; 87%). <sup>1</sup>H NMR (600 MHz, DMSO-d<sub>6</sub>): 0.70 and 0.73 (each d, J= 6.6 Hz, 6 H, 2 x Me), 0.85 and 1.55 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.15 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 2.54 (d, J= 4.4 Hz, 3 H, CONHCH<sub>3</sub>), 2.55 (m, 1 H, CH-iBu), 2.80 and 2.92 (each m, 2 H, Phe methylene), 2.90 and 3.40 (each m, 2 H, CH<sub>2</sub>OH), 4.45 (m, 2  
20 H, CONHCH and CH<sub>2</sub>OH), 4.95 (s, 1 H, tertiary OH), 7.1-7.3 (m, 5 H, Ph), 7.83 (broad s, 1 H, CONHCH<sub>3</sub>), 8.12 (broas s, 1 H, CONHCH), 8.65 (broad s, 1 H, NHOH), and 10.14 ppm (broad s, 1  
25 H, NHOH).

## EXAMPLE 8

**[(2R,3S)-4-Hydroxyamino-2-isobutyl-3-(2-tetrahydropyranyl)oxy-3-(2-tetrahydropyranyl)oxymethylsuccinyl]-L-tert-leucine-N-methylamide**



5

**Step 1: [(2R,3S)-4-Benzhydryloxy-2-isobutyl-3-(2-tetrahydropyranyl)oxy-3-(2-tetrahydropyranyl)oxymethyl-succinyl]-L-tert-leucine-N-methylamide**

A solution of the compound from Example 1, Step 3 2.7 g, 5.3 mmol) in dichloromethane (50 mL) was treated at 0°C with pyridinium-p-toluenesulfonate (150 mg) and 3,4-dihydro-2H-pyran (10 mL, 11 mmol). The mixture was kept in a cold room (4°C) for 3 days, then diluted with ethyl acetate (150 mL) and sequentially washed with 4% aqueous HCl, saturated aqueous sodium bicarbonate, and water. After drying over Na<sub>2</sub>SO<sub>4</sub> and evaporation, a colourless oil was obtained, which was treated with petroleum ether (150 mL) and kept at 4°C overnight. Fractionation by silica gel chromatography from the mono-pyranyl adduct (Example 1, Step 4) afforded the title compound.

20

**Step 2: [(2R,3S)-4-Hydroxy-2-isobutyl-3-(2-tetrahydropyranyl)oxy-3-(2-tetrahydropyranyl)oxymethylsuccinyl]-L-tert-leucine-N-methylamide**

The compound from Step 1 above (0.9 g) was taken up in ethanol (35 mL) and exposed to a hydrogen atmosphere in the presence of 10% Pd on charcoal (0.43 g) overnight. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol, and the combined ethanol solution was evaporated to dryness. The residue was taken up with n-hexane, stirred for 30 min, and the resulting white powder was collected by filtration, thereby obtaining the title compound (620 mg).

10 **Step 3: [(2R,3S)-2-Isobutyl-4-(4-nitrobenzyloxyamino)-3-(2-tetrahydropyranyl)oxy-3-(2-tetrahydropyranyl)oxymethylsuccinyl]-L-tert-leucine-N-methylamide**

A solution of the compound from Step 2 above (620 mg) in acetonitrile (35 mL) was treated with O-(4-nitrobenzyl)-hydroxylamine hydrochloride (324 mg), N-methylmorpholine (0.35 mL) and TBTU (555 mg) for 20 hr at room temperature. The mixture was taken up in ethyl acetate, and sequentially washed with 4% HCl and 4% sodium hydrogen carbonate. Evaporation and flash-chromatography over silica (EtOAc/hexane) afforded the title compound as a white solid (385 mg).


20 **Step 4: [(2R,3S)-4-Hydroxyamino-2-isobutyl-3-(2-tetrahydropyranyl)oxy-3-(2-tetrahydropyranyl)oxymethylsuccinyl]-L-tert-leucine-N-methylamide**

The compound from Step 3 above (100 mg) was taken up in ethanol (15 mL) and exposed to a hydrogen atmosphere in the presence of 10% Pd on charcoal (20 mg) for 12 hr. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol, and the combined ethanol solution was evaporated to dryness. The residue was taken up with n-hexane, stirred for 30 min, and the resulting white powder was collected by filtration, thereby obtaining the title compound

(diastereomeric mixture at the anomeric tetrahydropyran carbon atom; 27 mg).  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO-d}_6$ ): 0.75 (d,  $J = 6.8$  Hz, 6 H, 2 x Me), 0.87 (s, 9 H, t-Bu), 1.0-1.8 (m, 15 H,  $\text{CH}_2\text{CHMe}_2$  and pyran 3,4,5-methylene), 2.51 (d,  $J = 4.7$  Hz, 3 H,  $\text{CONHCH}_3$ ),  
 5 2.80 (m, 1 H,  $\text{CH-iBu}$ ), 3.3-3.9 (m, 6 H,  $\text{CH}_2\text{O-pyran}$  and pyran 6-methylene), 4.03 (d,  $J = 9.0$  Hz, 1 H,  $\text{CONHCH}$ ), 4.46 and 5.20 (each m, 1 H, anomeric CH of two pyrans), 7.40 (broad s, 1 H,  $\text{CONHCH}_3$ ), 7.70 (m, 1 H,  $\text{CONHCH}$ ), 8.78 (broad s, 1 H,  $\text{NHOH}$ ), and 10.31 ppm (broad s, 1 H,  $\text{NHOH}$ ).

#### EXAMPLE 9

***[(2R,3S)-3-Acetoxymethyl-3-hydroxy-4-hydroxyamino-2-isobutyl-succinyl]-L-tert-leucine-N-methylamide***



***Step 1: [(2R,3S)-3-Acetoxymethyl-4-benzyloxy-3-hydroxy-2-isobutylsuccinyl]-L-tert-leucine-N-methylamide***

A crude sample of (2R,3S)-(3-acetoxymethyl-4-benzyloxy-3-hydroxy-2-isobutyl)succinic acid (1.77 g, 5.02 mmol), obtained as described in Preparation 4, was dissolved in dry DMF (10 mL). To this solution, L-tert-leucine-N-methylamide (0.72 g, 5.02 mmol) and HOBT (1-hydroxybenzotriazole hydrate; 658 mg, 4.87 mmol) were added at room temperature, followed by dropwise addition at 0°C of a solution of DCC (dicyclohexylcarbodiimide; 1.04 g, 5.02 mmol) in THF (10 mL). The mixture was kept at 0°C for 30 min, then kept under

stirring for 5 h at room temperature, and finally allowed to stand overnight at 0°C. Dicyclohexylurea was filtered off, the solvent was removed in vacuo, the residue was taken up in ethyl acetate, and the solution was sequentially washed with  
5 10% aqueous citric acid (50 mL), saturated aqueous sodium bicarbonate (50 mL), and brine (50 mL). After drying over Na<sub>2</sub>SO<sub>4</sub>, the solvent was removed to give an oil. This residue was taken up in ethyl acetate and left aside to allow for separation of a further amount of dicyclohexylurea. After  
10 filtration and evaporation, the residue was triturated with isopropyl ether, thereby obtaining a first crop of the title product (0.95 g). Examination of the mother liquors revealed the presence of the benzotriazolyl ester ("active ester") of the starting material. After evaporation, this material was  
15 allowed to react with a further amount of L-tert-butyl-glycine-N-methylamide (0.3 g) in dry DMF (5 mL) for 65 hr at room temperature. Work-up as before gave a second crop of the title product, crystallizing from ethyl ether (280 mg). Flash chromatography over silica of the mother liquors gave a third  
20 crop (230 mg). The three crops were combined (1.46 g, 3.06 mmol; 61%).

**Step 2: [(2R,3S)-3-Acetoxymethyl-3,4-dihydroxy-2-isobutyl-succinyl]-L-tert-leucine-N-methylamide**

The compound from Step 1 above (0.9 g, 1.88 mmol) was taken up  
25 in ethanol (50 mL) and exposed to a hydrogen atmosphere in the presence of 5% Pd on charcoal (0.35 g) for 7 hr. A further amount of the catalyst (0.15 g) was added, and hydrogenation was continued for 16 hr. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional  
30 ethanol, and the combined ethanol solution was evaporated to dryness, thereby obtaining the title product (essentially pure



by HPLC) as a white solid (0.73 g, 1.88 mmol; quantitative yield).

**Step 3: [(2R,3S)-3-Acetoxymethyl-4-benzyloxyamino-3-hydroxy-2-isobutylsuccinyl]-L-tert-leucine-N-methylamide**

5 The compound from Step 2 above (720 mg, 1.86 mmol) was dissolved in acetonitrile (25 mL). To this solution, O-benzylhydroxylamine hydrochloride (326 mg, 2.05 mmol), N-methylmorpholine (0.45 mL, 4.09 mmol), and O-benzotriazol-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU; 717 mg, 10 2.23 mmol) were added in this order. After stirring for 26 h at room temperature, the solvent was removed under reduced pressure and the residue was taken up in ethyl acetate. Sequential washing with saturated aqueous sodium bicarbonate, 2N aqueous HCl and water, followed by drying over Na<sub>2</sub>SO<sub>4</sub> and 15 evaporation, left a residue which was purified by flash chromatography over silica (n-hexane and ethyl acetate gradient elution), thereby obtaining the title product (230 mg, 0.46 mmol; 25%).

**Step 4: [(2R,3S)-3-Acetoxymethyl-3-hydroxy-4-hydroxyamino-2-isobutylsuccinyl]-L-tert-leucine-N-methylamide**


20 A solution of the compound from Step 3 above (230 mg, 0.46 mmol) in ethanol (20 mL) was exposed to a hydrogen atmosphere in the presence of 5% Pd on charcoal (74 mg) for 90 min. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), 25 washed with additional ethanol, and the combined ethanol solution was evaporated to dryness. The residue was taken up with isopropyl ether, stirred for 15 min, and the resulting white powder was collected by filtration, thereby obtaining the title compound as a white solid (135 mg, 0.335 mmol; 73%). 30 <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.77 and 0.80 (each d, J= 6.7 Hz, 6 H, 2 x Me), 0.91 (s, 9 H, t-Bu), 0.96 and 1.63 (each m, 2 H,

72

CH<sub>2</sub>CHMe<sub>2</sub>), 1.28 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.94 (s, 3 H, COCH<sub>3</sub>), 2.54 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>), 2.97 (dd, J= 3.5 and 11.7 Hz, 1 H, CH-iBu), 3.99 (d, J= 10.0 Hz, 1 H, CONHCH), 4.15 (m, 2 H, CH<sub>2</sub>OAc), 5.40 (s, 1 H, tertiary OH), 7.88 (q, J= 4.7 Hz, 1 H, CONHCH<sub>3</sub>), 8.19 (d, J= 10.0 Hz, 1 H, CONHCH), 8.80 (s, 1 H, NHOH), and 10.42 ppm (broad s, 1 H, NHOH).

**EXAMPLE 10**

**[(2R,3S)-3-Acetoxymethyl-3-hydroxy-4-hydroxyamino-2-isobutyl-succinyl]-L-phenylalanine-N-methylamide**



10


Starting from (2R,3S)-(3-Acetoxymethyl-4-benzyloxy-3-hydroxy-2-isobutyl)succinic acid (Preparation 4), but substituting L-phenylalanine-N-methylamide for L-tert-leucine-N-methylamide, and following a procedure analogous to that described in Example 9, the title product was obtained. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.70 and 0.720 (each d, J= 6.5 Hz, 6 H, 2 x Me), 0.80 and 1.70 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.20 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.88 (s, 3 H, COCH<sub>3</sub>), 2.50 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>), 2.62 (m, 1 H, CH-iBu), 2.77 and 2.89 (each m, 1 H, CH<sub>2</sub>Ph), 3.48 and 3.95 (each d, J= 11.1 Hz, CH<sub>2</sub>OAc), 4.48 (m, 1 H, CONHCH), 5.20 (s, 1 H, tertiary OH), 7.0-7.2 (m, 5 H, Ph), 7.82 (broad s, 1 H, CONHCH<sub>3</sub>), 8.42 (broad s, 1 H, CONHCH), 8.65 (broad s, 1 H, NHOH), and 10.37 ppm (broad s, 1 H, NHOH).

25

73

## EXAMPLE 11


[(2R,3S)-3-Acetoxymethyl-3-hydroxy-4-hydroxyamino-2-isobutyl-succinyl]-L-tert-leucine-N-(4-pyridyl)amide



Starting from (2R,3S)-(3-Acetoxymethyl-4-benzyloxy-3-hydroxy-2-isobutyl)succinic acid (Preparation 4), but substituting L-phenylalanine-N-methylamide for L-tert-leucine-N-(4-pyridyl)-amide, and following a procedure analogous to that described in Example 9, the title product was obtained. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.73 and 0.80 (each d, J= 6.4 Hz, 6 H, 2 x Me), 0.90 and 1.65 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 0.99 (s, 9 H, t-Bu), 1.30 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.94 (s, 3 H, COCH<sub>3</sub>), 3.05 (dd, J= 3.6 and 11.5 Hz, 1 H, CH-iBu), 4.03 and 4.17 (each d, J= 11.1 Hz, 1 H, CH<sub>2</sub>OAc), 4.35 (d, J= 8.1 Hz, 1 H, CONHCH), 5.24 (s, 1 H, tertiary OH), 7.52 (d, J= 6.4 Hz, 2 H, 2,6-pyridyl protons), 8.40 (d, J= 6.4 Hz, 2 H, 3,5-pyridyl protons), 8.40 (s, 1 H, CONHCH), 8.82 (broad s, 1 H, NHOH), 10.48 (s, 1 H, CONH-pyr), and 10.50 ppm (broad s, 1 H, NHOH).

## EXAMPLE 12

[(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutyl-succinyl]-L-tert-leucine-N-(4-pyridyl)amide




A solution of [(2R,3S)-3-Acetoxymethyl-3-hydroxy-4-hydroxy-amino-2-isobutyl-succinyl]-L-tert-leucine-N-(4-pyridyl)amide  
 5 (Example 11; 10.5 mg) in methanol (2 mL) was treated with sodium methoxide (2.45 mg) for 1 hr at room temperature. After addition of acetic acid (0.025 mL), the solution was passed through a reverse-phase (RP-C-18) column, eluting with EtOH/H<sub>2</sub>O 3:1. Pooled fractions were lyophilized, to afford the title  
 10 compound as a white lyophile, 8 mg. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.74 and 0.80 (each d, J= 6.4 Hz, 6 H, 2 x Me), 0.90 and 1.65 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 0.99 (s, 9 H, t-Bu), 1.30 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 2.90 (dd, J= 3.6 and 11.7 Hz, 1 H, CH-iBu), 3.20 and 3.65 (each m, 1 H, CH<sub>2</sub>OH), 4.32 (d, J= 8.2 Hz, 1 H, CONHCH), 4.68 (dd, J= 4.4 and 7.0 Hz, CH<sub>2</sub>OH), 5.04 (s, 1 H, tertiary OH), 7.53 (d, J= 6.1 Hz, 2 H, 2,6-pyridyl protons), 8.12 (d, J= 8.2 Hz, 1 H, CONHCH), 8.40 (d, J= 6.1 Hz, 2 H, 3,5-pyridyl protons), 8.72 (d, J= 2.0 Hz, 1 H, NHOH), 10.22 (d, J= 2.0 Hz, 1 H, NHOH), and 10.45 ppm (s, 1 H, CONH-pyr).

20

**EXAMPLE 13**

**[(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(trimethyl-acetoxy)methyl-succinyl]-L-tert-leucine-N-methylamide**



**Step 1: (2R,3S)-[4-Benzyloxy-3-hydroxy-2-isobutyl-3-(trimethyl-acetoxy)methyl]succinic acid**

A solution in dichloromethane (40 mL) of tert-butyl (2R,3S)-  
 5 (4-benzyloxy-3-hydroxy-3-hydroxymethyl-2-isobutyl)succinate  
 (1.83 g), obtained as described in Preparation 3, was treated  
 at 0 °C with pivaloyl chloride (0.74 mL) and triethylamine  
 (0.84 mL). After standing for 48 hr at room temperature, the  
 solution was sequentially washed with water, 4% HCl, and 4%  
 10 sodium hydrogen carbonate. Evaporation in vacuo left a  
 residue, which was taken up in dichloromethane (5 mL) and  
 treated at room temperature for 2 hr with trifluoroacetic acid  
 (4 mL) and anisole (0.5 mL). Concentration and flash-  
 chromatography gave the title product as a waxy solid (1.55  
 15 g).

**Step 2: [(2R,3S)-4-Benzyloxy-3-hydroxy-2-isobutyl-3-(trimethyl-acetoxy)methyl-succinyl]-L-tert-leucine-N-methylamide**

A solution of the compound from Step 1 above (925 mg) in dry  
 20 DMF (5 mL) was treated with L-tert-leucine-N-methylamide (375  
 mg) and triethylamine (0.3 mL). After cooling to 0 °C, a  
 solution of DCC (540 mg) in dry DMF (3 mL) was added. The  
 mixture was let rise to room temperature, left aside for 6  
 days, and finally stirred at 50 °C for 24 hr. Partitioning

between EtOAc and 4% HCl, followed by washing with 4% sodium hydrogen carbonate and evaporation of the solvent, afforded the crude title product (800 mg).

**Step 3: [(2R,3S)-3,4-Dihydroxy-2-isobutyl-3-(trimethylacetoxymethylsuccinyl)-L-tert-leucine-N-methylamide]**

The compound from Step 2 above (700 mg) in ethanol (50 mL) was exposed to a hydrogen atmosphere in the presence of 5% Pd/C (330 mg) at room temperature for 30 hr. Filtration of the catalyst and evaporation in vacuo left the crude title product (465 mg).

**Step 4: [(2R,3S)-4-Benzoyloxyamino-3-hydroxy-2-isobutyl-3-(trimethylacetoxymethylsuccinyl)-L-tert-leucine-N-methylamide]**

The compound from Step 3 above (450 mg) in acetonitrile (30 mL) was treated with O-(benzyl)hydroxylamine hydrochloride (188 mg), N-methylmorpholine (0.258 mL), and TBTU (412 mg) at room temperature for 18 hr. Work-up (EtOAc/water, 4% HCl, saturated sodium hydrogen carbonate) and flash-chromatography afforded the title product as a white solid (103 mg).

**Step 5: [(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(trimethylacetoxymethylsuccinyl)-L-tert-leucine-N-methylamide]**


The compound from Step 4 above (100 mg) in ethanol (10 mL) was exposed to a hydrogen atmosphere in the presence of 5% Pd/C (35 mg) at room temperature for 2 hr. Filtration of the catalyst and evaporation in vacuo afforded the title product as a white solid (77 mg). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.73 and 0.77 (each d, J= 6.4 Hz, 6 H, 2 x Me), 0.88 (s, 9 H, t-Bu of t-leucine), 1.05 (s, 9 H, t-Bu of pivalate), 0.90 and 1.60 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.23 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 2.50 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>), 2.98 (dd, J= 3.4 and 11.9 Hz, 1 H, CH-

77

iBu), 3.98 and 4.09 (each d,  $J = 11.1$  Hz, 2 H,  $\text{CH}_2\text{OCO}$ ), 4.12 (d,  $J = 8.9$  Hz, 1 H,  $\text{CONHCH}$ ), 5.26 (s, 1 H, tertiary OH), 7.85 (q,  $J = 4.7$  Hz, 1 H,  $\text{CONHCH}_3$ ), 8.20 (d,  $J = 8.9$  Hz, 1 H,  $\text{CONHCH}$ ), 8.72 (s, 1 H,  $\text{NHOH}$ ), and 10.35 ppm (broad s, 1 H,  $\text{NHOH}$ ).

**EXAMPLE 14**

**[(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-tert-butylamide**



**Step 1: [(2R,3S)-3-Acetoxymethyl-4-benzyloxy-3-hydroxy-2-isobutylsuccinyl]-L-tert-leucine-N-tert-butylamide**  
 (2R,3S)-(3-Acetoxymethyl-3-benzyloxycarbonyl-2-isobutyl)-propiolactone (1.6 g), obtained as described in Preparation 9, was dissolved in dry DMF (10 mL). To this solution, L-tert-leucine-N-tert-butylamide (0.85 g) and a catalytic amount of HOBT (1-hydroxybenzotriazole hydrate; 80 mg) were added at room temperature. After stirring overnight at room temperature, the solvent was removed in vacuo, the residue was taken up in ethyl acetate, and the solution was sequentially washed with 10% aqueous citric acid (50 mL), saturated aqueous sodium bicarbonate (50 mL), and brine (50 mL). Drying over  $\text{Na}_2\text{SO}_4$  and evaporation in vacuo afforded the crude title product as a syrup.

**Step 2: [(2R,3S)-3-Acetoxymethyl-3,4-dihydroxy-2-isobutylsuccinyl]-L-tert-leucine-N-tert-butylamide**

The crude compound from Step 1 above was taken up in ethanol (100 mL) and exposed to a hydrogen atmosphere in the presence

78

of 5% Pd on charcoal (0.7 g) for 7 hr. A further amount of the catalyst (0.3 g) was added, and hydrogenation was continued for 16 hr. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol, and the combined ethanol solution was evaporated to dryness, thereby obtaining the title product (1.5 g).

**Step 3: [(2R,3S)-3-Acetoxymethyl-4-benzyloxyamino-3-hydroxy-2-isobutylsuccinyl]-L-tert-leucine-N-tert-butylamide**

The compound from Step 2 above was dissolved in acetonitrile (50 mL). To this solution, O-benzylhydroxylamine hydrochloride (660 mg), N-methylmorpholine (0.9 mL), and O-benzotriazol-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU; 1.5 g) were added in this order. After stirring for 24 h at room temperature, the solvent was removed under reduced pressure and the residue was taken up in ethyl acetate. Sequential washing with saturated aqueous sodium bicarbonate, 2N aqueous HCl and water, followed by drying over Na<sub>2</sub>SO<sub>4</sub> and evaporation, left a residue which was purified by flash chromatography over silica (n-hexane and ethyl acetate gradient elution), thereby obtaining the title product (0.7 g).

**Step 4: [(2R,3S)-3-Acetoxymethyl-3-hydroxy-4-hydroxyamino-2-isobutylsuccinyl]-L-tert-leucine-N-tert-butylamide**

A solution of the compound from Step 3 above in ethanol (60 mL) was exposed to a hydrogen atmosphere in the presence of 5% Pd on charcoal (200 mg) for 90 min. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol, and the combined ethanol solution was evaporated to dryness. The residue was taken up with isopropyl ether, stirred for 15 min, and the resulting white powder was collected by filtration, thereby obtaining the title compound as a white solid (400 mg).




**Step 5: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutyl-succinyl]-L-tert-leucine-N-tert-butylamide**

A solution of the compound from Step 4 above in methanol (50 mL) was treated with sodium methoxide (100 mg) for 1 hr at room temperature. After addition of acetic acid (0.1 mL), the solution was passed through a reverse-phase (RP-C-18) column, eluting with EtOH/H<sub>2</sub>O 3:1. Pooled fractions were lyophilized, to afford the title compound as a white lyophile, 320 mg. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.76 and 0.79 (each d, J= 6.8 Hz, 6 H, 2 x Me), 0.90 (s, 9 H, t-Bu of t-leucine), 1.21 (s, 9 H, CONH-t-Bu), 0.90 and 1.65 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.30 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 2.85 (dd, J= 3.4 and 11.9 Hz, 1 H, CH-iBu), 3.22 (dd, J= 4.7 and 10.7 Hz, 1 H, CH(H)OH), 3.63 (dd, J= 7.3 and 10.7 Hz, 1 H, CH(H)OH), 4.18 (d, J= 9.8 Hz, 1 H, CONHCH), 4.59 (dd, J= 4.7 and 7.3 Hz, 1 H, CH<sub>2</sub>OH), 5.14 (s, 1 H, tertiary OH), 7.50 (s, 1 H, CONH-tBu), 7.94 (d, J= 9.8 Hz, 1 H, CONHCH), 8.68 (d, J= 2.1 Hz, 1 H, NHOH), and 10.17 ppm (d, J= 2.1 Hz, 1 H, NHOH).

**EXAMPLE 15**

**[(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutyl-succinyl]-L-cyclohexylglycine-N-tert-butylamide**




Starting from (2R,3S)-(3-acetoxymethyl-3-benzyloxycarbonyl-2-isobutyl)-propiolactone, obtained as described in Preparation 9, and by the same procedure as that described in Example 14, but replacing L-tert-leucine-N-tert-butylamide with L-

80

cyclohexylglycine-N-tert-butylamide, the title compound was obtained.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ ): 0.76 and 0.79 (each d,  $J=6.4$  Hz, 6 H, 2 x Me), 1.20 (s, 9 H, CONH-t-Bu), 0.8-1.7 (m, 14 H,  $\text{CH}_2\text{CHMe}_2$  and cyclohexyl protons), 2.73 (dd,  $J=3.4$  and 11.9 Hz, 1 H,  $\text{CH-iBu}$ ), 3.20 (dd,  $J=4.7$  and 10.8 Hz, 1 H,  $\text{CH(H)OH}$ ), 3.59 (dd,  $J=7.3$  and 10.8 Hz, 1 H,  $\text{CH(H)OH}$ ), 4.05 (d,  $J=8.5$  Hz, 1 H, CONHCH), 4.60 (dd,  $J=4.7$  and 7.3 Hz, 1 H,  $\text{CH}_2\text{OH}$ ), 5.12 (s, 1 H, tertiary OH), 7.38 (s, 1 H, CONH-tBu), 8.16 (d,  $J=8.5$  Hz, 1 H, CONHCH), 8.67 (broad s, 1 H,  $\text{NHOH}$ ), and 10.17 ppm (broad s, 1 H,  $\text{NHOH}$ ).

**EXAMPLE 16**

***[(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-3-hydroxy-4-hydroxyamino-2-isobutylsuccinyl]-L-tert-leucine-N-methylamide***



**Step 1: (2R,3S)-4-Benzylloxy-3-(N-cyclohexylaminocarbonyl)-oxymethyl-3-hydroxy-2-isobutylsuccinic acid**

4-Dimethylaminopyridine (200 mg) and cyclohexylisocyanate (5 mL) were added to a solution in dichloromethane (30 mL) of tert-butyl (2R,3S)-(4-benzylloxy-3-hydroxy--3-hydroxymethyl -2-isobutyl) succinate (6.5 g), obtained as described in Preparation 3. After stirring for 16 hr at room temperature, the reaction mixture was quenched with methanol (20 mL) and the solvent removed in vacuo. Cold (-20 °C) 95% aqueous trifluoroacetic acid (30 mL) was added to the residue, and the solution was let standing for 4 hr at room temperature.

Concentration and flash-chromatography over silica (hexane/  
EtOAc gradient elution, from 80:20 to 0:100) and evaporation  
of pooled fractions left a waxy residue, which was triturated  
with isopropyl ether to give the title product as a white  
5 powder (5.9 g).

**Step 2: [(2R,3S)-4-Benzoyloxy-3-(N-cyclohexylaminocarbonyl)-  
oxymethyl-3-hydroxy-2-isobutylsuccinyl]-L-tert-leucine-N-  
methylamide**

The compound from Step 1 above (5.9 g) was dissolved in a  
10 mixture of DMF (45 mL) and dichloromethane (30 mL). After  
cooling to 0 °C, HOBT (2.03 g), N,N-dimethylaminopropyl-N'-  
ethyl carbodiimide hydrochloride (2.88 g) and N-  
methylmorpholine (1.76 mL) were added, followed, after 1 hr  
at 0 °C, by tert-leucine-N-methylamide (2.88 g). The reaction  
15 was let standing for 2 days at room temperature, then heated  
at 65 °C for 3 hr. Partitioning between ethyl acetate and  
water, followed by sequential washing with 4% HCl and  
saturated sodium hydrogen carbonate, drying (Na<sub>2</sub>SO<sub>4</sub>) and  
evaporation, left the crude title product. Flash-  
20 chromatography over silica afforded the pure compound as a  
white foam, 6.3 g.

**Step 3: [(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-3,4-  
dihydroxy-2-isobutylsuccinyl]-L-tert-leucine-N-methylamide**

The compound from Step 2 above (6.3 g) in ethanol (400 mL) was  
25 exposed to a hydrogen atmosphere for 2 hr in the presence of  
5% Pd/C (1.5 g). Filtration of the catalyst and evaporation  
gave the title compound as a white foam (5.1 g).

**Step 4: [(2R,3S)-4-Benzoyloxyamino-3-(N-cyclohexylamino-  
carbonyl)oxymethyl-3-hydroxy-2-isobutylsuccinyl]-L-tert-  
30 leucine-N-methylamide**

The compound from Step 3 above (4.15 g) in acetonitrile (150 mL) was treated with O-(benzyl)hydroxylamine hydrochloride (1.67 g), N-methylmorpholine (2.14 mL) and TBTU (3.37 g). After 20 hr at room temperature, aqueous work-up and flash-  
5 chromatography gave the title product as a white foam (1.8 g).

**Step 5: [(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-3-hydroxy-4-hydroxyamino-2-isobutylsuccinyl]-L-tert-leucine-N-methylamide**

The compound from Step 4 above (1.8 g) in ethanol (150 mL) was  
10 exposed to a hydrogen atmosphere for 5 hr in the presence of 5% Pd/C (150 mg). Filtration of the catalyst and evaporation left a waxy solid, which was triturated with ethyl ether (3 x 50 mL) to afford the title compound as a white powder (1.25 g).  
<sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.76 and 0.80 (each d, J= 6.4 Hz, 6  
15 H, 2 x Me), 0.90 (s, 9 H, t-Bu), 0.9-1.7 (m, 13 H, CH<sub>2</sub>CHMe<sub>2</sub> and cyclohexyl methylene protons), 2.97 (m, 1 H, CH-iBu), 3.17 (m, 1 H, cyclohexyl methine proton), 3.91 and 4.06 (each d, J= 11.7 Hz, 2 H, CH<sub>2</sub>OCO), 4.14 (d, J= 9.4 Hz, 1 H, CONHCH), 5.32 (s, 1 H, tertiary OH), 6.89 (d, J= 7.7 Hz, 1 H, CONH-  
20 cyclohexyl), 7.87 (q, J= 4.3 Hz, 1 H, CONH-Me), 8.12 (d, J= 9.4 Hz, 1 H, CONHCH), 8.75 (broad s, 1 H, NHOH), and 10.34 ppm (broad s, 1 H, NHOH).

**EXAMPLE 17**

By procedures similar to those described in Examples 1-16  
25 above, the following compounds may be prepared:

#17.1: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-phenylalanine-N-tert-butylamide;

#17.2: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-phenylalanine-N-2-(morpholino)ethylamide;

30 #17.3: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-(O-tert-butyl)threonine-N-methylamide;

- #17.4: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-amide;
- #17.5: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-cyclopropylamide;
- 5 #17.6: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-cyclopentylamide;
- #17.7: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-(3-quinolyl)amide;
- #17.8: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-
- 10 isobutylsuccinyl]-L-tert-leucine-N-(5-quinolyl)amide;
- #17.9: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-(3-isoquinolyl)amide;
- #17.10: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-(3-quinuclidinyl)amide;
- 15 #17.11: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-(3,4-methylenedioxy)anilide;
- #17.12: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-(S-methyl)penicillamine-N-methylamide;
- 20 #17.13: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-[(S-methyl)penicillamine-sulfoxide]-N-methylamide;
- #17.14: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-[(S-methyl)penicillamine-sulfone]-N-
- 25 methylamide;
- #17.15: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-cyclohexylalanine-N-methylamide;
- #17.16: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-cyclohexylalanine-N-tert-butylamide;

- #17.17: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-cyclohexylalanine-N-2-(1-morpholino)-ethylamide;
- #17.18: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-cyclohexylalanine-N-2-(4-aminosulfonyl-phenyl)ethylamide;
- #17.19: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-phenylalanine-N-2-(4-aminosulfonyl-phenyl)ethylamide;
- #17.20: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-2-(4-aminosulfonyl-phenyl)ethylamide;
- #17.21: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-(4-piperidyl)amide;
- #17.22: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-cyclohexylglycine-N-(4-piperidyl)amide;
- #17.23: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-cyclohexylglycine-N-(4-pyridyl)amide;
- #17.24: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-2-pyridylalanine-N-tert-butylamide;
- #17.25: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-3-pyridylalanine-N-(3,4-dioxymethylene)-anilide;
- #17.26: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-7-isoquinolylalanine-N-methylamide;
- #17.27: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-2-quinolylalanine-N-methylamide;
- #17.28: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-(4-fluoro)phenylalanine-N-methylamide;

- #17.29: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-(4-fluoro)phenylalanine-N-(4-pyridyl) amide;
- #17.30: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-(4-thiazolyl) alanine-N-(4-pyridyl) amide;
- 5 #17.31: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-(4-thiazolyl) alanine-N-tert-butylamide;
- #17.32: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-(5,6,7,8-tetrahydro-7-isoquinolyl) alanine-
- 10 N-methylamide;
- #17.33: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-(1-adamantyl) glycine-N-methylamide;
- #17.34: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-(2-methoxyethyl) alanine-N-tert-butylamide;
- 15 #17.35: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-[1-phenyl-1-(3-pyridyl)]-methylamide;
- #17.36: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-tert-leucine-N-methylamide;
- 20 #17.37: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-tert-leucine-N-tert-butylamide;
- #17.38: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-tert-leucine-carboxamide;
- #17.39: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-
- 25 3-hydroxymethylsuccinyl]-L-tert-leucine-N-(4-pyridyl) amide;
- #17.40: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-(S-methyl) penicillamine-N-methylamide;
- #17.41: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-
- 30 3-hydroxymethylsuccinyl]-L-cyclohexylalanine-N-methylamide;

- #17.42: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-cyclohexylalanine-N-2-(1-morpholino)-ethylamide;
- #17.43: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-cyclohexylalanine-N-2-(4-amino-sulfonylphenyl)ethylamide;
- #17.44: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-tert-leucine-N-2-(4-aminosulfonylphenyl)ethylamide;
- #17.45: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-cyclohexylglycine-N-(4-pyridyl)amide;
- #17.46: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-3-pyridylalanine-N-(3,4-dioxy-methylene)anilide;
- #17.47: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-(4-fluoro)phenylalanine-N-methylamide;
- #17.48: [(2R,3S)-2-(3-(4-Chlorophenyl)propyl)-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-tert-leucine-N-methylamide;
- #17.49: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-(3-(4-methoxyphenyl)propyl)succinyl]-L-tert-leucine-N-methylamide;
- #17.50: [(2R,3S)-2-(3-(4-Diphenyl)propyl)-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-tert-leucine-N-(4-pyridyl)amide;
- #17.51: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-(3-(4-methoxyphenyl)oxypropyl)succinyl]-L-tert-leucine-N-methylamide;



- #17.52: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-(5-phenylpentyl)succinyl]-L-tert-leucine-N-methylamide;
- #17.53: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-(4-phenoxybutyl)succinyl]-L-tert-leucine-N-methylamide;
- 5 #17.54: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-(5-(4-propylphenyl)oxypentyl)succinyl]-L-tert-leucine-amide;
- #17.55: [(2R,3S)-2-(3-(Benzylaminocarbonyl)propyl)-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-tert-leucine-N-methyl-amide;
- 10 #17.56: [(2R,3S)-2-(3-(3,4-Difluorophenyl)aminocarbonyl)-propyl)-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-tert-leucine-N-methylamide;
- #17.57: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-octylsuccinyl]-L-tert-leucine-N-methylamide;
- 15 #17.58: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-octylsuccinyl]-L-tert-leucine-N-(4-pyridyl)amide;
- #17.59: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-(4-methoxyphenylsulfonyl)succinyl]-L-tert-leucine-N-methylamide;
- 20 #17.60: [(2R,3S)-2-(4-Butoxyphenylsulfonyl)-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-L-tert-leucine-N-methylamide;
- #17.61: [(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-2-cyclopentylmethyl-3-hydroxy-4-hydroxyaminosuccinyl]-L-tert-
- 25 leucine-N-methylamide;
- #17.62: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-(N-piperidyl)carbonyloxymethyl-succinyl]-L-tert-leucine-N-methylamide;
- #17.63: [(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-
- 30 phenoxyethylsuccinyl]-L-tert-leucine-N-methylamide;

#17.64: [(2R,3S)-3-(4-Aminophenyl)oxymethyl-3-hydroxy-4-hydroxyamino-2-isobutylsuccinyl]-L-tert-leucine-N-methylamide;

#17.65: [(2R,3S)-2-Cyclopentylmethyl-3-(4-fluorophenyl)-oxymethyl-3-hydroxy-4-hydroxyaminosuccinyl]-L-tert-leucine-N-methylamide;

#17.66: [(2R,3S)-2-Cyclopentylmethyl-3-(3,4-dioxymethylene)-phenoxy-methyl-3-hydroxy-4-hydroxyaminosuccinyl]-L-tert-leucine-N-methylamide;


#17.67: [(2R,3S)-3-Hydroxy-4-hydroxyamino-3-(4-hydroxymethylphenyl)oxymethyl-2-isobutylsuccinyl]-L-tert-leucine-N-methylamide;

#17.68: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-methoxymethylsuccinyl]-L-tert-leucine-N-methylamide;

#17.69: [(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(3-pyridyl)oxymethylsuccinyl]-L-tert-leucine-N-methylamide.

#### EXAMPLE 18

[(2R,3S)-3,4-Dihydroxy-2-isobutyl-3-(1-methyl-1,2,3,4-tetrazol-5-yl)thiomethylsuccinyl]-L-phenylalanine-N-methylamide



20

Step 1: [(2R,3S)-4-Benzhydryloxy-3-hydroxy-2-isobutyl-3-(1-methyltetrazol-5-yl)thiomethylsuccinyl]-L-phenylalanine-N-methylamide, and

**[2R-[(2S)-2-Benzhydryloxy-carbonyl-2-oxiranyl]-4-methyl-pentanoyl]-L-phenylalanine-N-methylamide**

A solution in dichloromethane (10 mL) of (2R,3S)-(4-benzhydryloxy-3-hydroxy-3-hydroxymethyl-2-isobutylsuccinyl)-L-phenylalanine-N-methylamide (100 mg; 0.183 mmol), prepared as described in Example 4, Step 3, was treated with triphenylphosphine (140 mg, 0.53 mmol), 5-mercapto-1-methyltetrazole (50 mg, 0.43 mmol), and diethyl azodicarboxylate (0.09 mL, 0.57 mmol). After stirring for 2 h at room temperature, the reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed, in this order, with 2% aqueous HCl, saturated aqueous sodium bicarbonate and brine, then dried over sodium sulfate and evaporated to dryness. Fractionation by flash chromatography over silica (n-hexane and ethyl acetate gradient elution) afforded, in this order:

[2R-[(2S)-2-Benzhydryloxy-carbonyl-2-oxiranyl]-4-methyl-pentanoyl]-L-phenylalanine-N-methylamide, white foam (30 mg, 0.056 mmol; 31%). IR (film),  $\nu_{\max}$  3296, 2957, 2870, 1736, and 1645  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ ): 0.71 and 0.72 (each d, J= 7.0 Hz, 6 H, 2 x Me), 1.05 and 1.55 (each m, 2 H,  $\text{CH}_2\text{CHMe}_2$ ), 1.30 (m, 1 H,  $\text{CHMe}_2$ ), 2.12 and 2.57 (each d, J= 5.6 Hz, 2 H, oxirane methylene), 2.54 (d, J= 4.4 Hz, 3 H,  $\text{CONHCH}_3$ ), 2.73-3.21 (m, 3 H, Phe methylene and  $\text{CH-iBu}$ ), 4.41 (m, 1 H,  $\text{CONHCH}$ ), 6.81 (s, 1 H,  $\text{CHPh}_2$ ), 7.0-7.6 (m, 15 H, 3 x Ph), 7.85 (q, J= 4.4 Hz, 1 H,  $\text{CONHCH}_3$ ), and 8.23 ppm (d, J= 8.5 Hz, 1 H,  $\text{CONHCH}$ ); and

[(2R,3S)-4-Benzhydryloxy-3-hydroxy-2-isobutyl-3-(1-methyl-tetrazol-5-yl)thiomethylsuccinyl]-L-phenylalanine-N-methylamide, (32 mg, 0.054 mmol; 30%), white foam, slightly impure from triphenylphosphine oxide.

**Step 2: [(2R,3S)-3,4-Dihydroxy-2-isobutyl-3-(1-methyltetrazol-5-yl)thiomethylsuccinyl]-L-phenylalanine-N-methylamide**

[(2R,3S)-4-Benzhydryloxy-3-hydroxy-2-isobutyl-3-(1-methyltetrazol-5-yl)thiomethylsuccinyl]-L-phenylalanine-N-

5 methylamide (20 mg, 0.031 mmol), as obtained from Step 1 above, was dissolved in dichloromethane (5 mL). Anisole (5 mL) and trifluoroacetic acid (1.5 mL) were added and the mixture left aside for 3 h. Concentration under reduced pressure left a waxy residue, which was triturated in ethyl ether. The title  
10 product was collected as a white solid by filtration (10 mg, 0.021 mmol; 68%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.68 and 0.70 (each d, J= 8.8 Hz, 6 H, 2 x Me), 0.76 and 1.62 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.14 (m, 1 H, CHMe<sub>2</sub>), 2.51 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>), 2.66 (dd, J= 3.0 and 12.0 Hz, 1 H, CH-iBu), 2.75 and  
15 2.92 (each m, 2 H, Phe methylene), 2.97 and 3.46 (each d, J= 13.2 Hz, 2 H, CH<sub>2</sub>S), 3.86 (s, 3 H, tetrazole N-CH<sub>3</sub>), 4.48 (m, 1 H, CONHCH), 5.38 (broad s, 1 H, tertiary OH), 7.0-7.2 (m, 5 H, Ph), 7.84 (q, J= 4.7 Hz, 1 H, CONHCH<sub>3</sub>), 8.19 (d, J= 8.1 Hz, 1 H, CONHCH), and 13.0 ppm (broad s, 1 H, COOH).

20

**EXAMPLE 19**

Similarly, the following compounds may be prepared:

#19.1: [(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(1-methyl-1,2,3,4-tetrazol-5-yl)thiomethylsuccinyl]-L-tert-leucine-N-methylamide;

25 #19.2: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-(2-thienyl)thiomethylsuccinyl]-L-tert-leucine-N-methylamide;

#19.3: [(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(2-pyridyl)thiomethylsuccinyl]-L-tert-leucine-N-methylamide;


#19.4: [(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(4-pyridyl)thiomethylsuccinyl]-L-tert-leucine-N-methylamide;

30

- #19.5: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-(4-hydroxyphenyl)thiomethyl-2-isobutylsuccinyl]-L-tert-leucine-N-methylamide;
- #19.6: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-methylthiomethylsuccinyl]-L-tert-leucine-N-methylamide;
- #19.7: [(2R,3S)-2-Cyclopentylmethyl-3-(4-fluorobenzene-sulfonyl)methyl-3-hydroxy-4-hydroxyaminosuccinyl]-L-tert-leucine-N-methylamide;
- #19.8: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-(4-methoxybenzenesulfonyl)methylsuccinyl]-L-tert-leucine-N-methylamide;
- #19.9: [(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-mercaptomethylsuccinyl]-L-tert-leucine-N-methylamide;
- #19.10: [(2R,3S)-3-Acetylthiomethyl-2-cyclopentylmethyl-3-hydroxy-4-hydroxyaminosuccinyl]-L-tert-leucine-N-methylamide.

## EXAMPLE 20

[2R-(1-Hydroxyaminocarbonyl)cyclopropyl-4-methylpentanoyl]-L-phenylalanine-N-methylamide



- Step 1: (4-Benzhydryloxy-2R-isobutyl-3-methylenesuccinyl)-L-phenylalanine-N-methylamide

(4-Hydroxy-2R-isobutyl-3-methylenesuccinyl)-L-phenylalanine-N-methylamide (2.0 g, 5.76 mmol), prepared as described in Example 4, Step 2, was dissolved in a mixture of acetonitrile (66 mL) and dichloromethane (34 mL). A solution of diphenyldiazomethane (DDM; 1.17 g, 6 mmol) in dichloromethane

(20 mL) was added dropwise under stirring, and the mixture left at room temperature for 1 h. Acetic acid (0.5 mL) was added to discharge the color of excess DDM. Evaporation under reduced pressure left a residue, which was treated with diisopropyl ether. The white precipitate was collected by filtration (2.5 g, 4.88 mmol; 85%). IR (KBr),  $\nu_{\max}$  3317, 3035, 2953, 2368, 1715, 1648, 1538 and 1455  $\text{cm}^{-1}$ .

**Step 2: [2R-(1-Benzhydryloxy carbonyl) cyclopropyl-4-methyl-pentanoyl]-L-phenylalanine-N-methylamide**

The compound from Step 1 above (500 mg, 0.976 mmol) was taken up in dichloromethane (2 mL) and ethyl ether (150 mL). To this solution, an ethereal solution of diazomethane (prepared from N-methyl-N-nitroso-p-toluenesulfonamide, 2 g, by Aldrich Diazald<sup>R</sup> Kit) was added dropwise. The mixture was left standing overnight, then evaporated and taken up in acetonitrile (150 mL). This solution was irradiated ( $\lambda > 300$  nm) until complete conversion into a new product was monitored by HPLC. The solvent was evaporated under reduced pressure and the residue was purified by flash chromatography (n-hexane and ethyl acetate gradient elution) to afford the title compound as a white foam (250 mg, 0.486 mmol; 50%). IR (KBr),  $\nu_{\max}$  3326, 3063, 3022, 2957, 1740, and 1652  $\text{cm}^{-1}$ .

**Step 3: [2R-(1-Carboxy) cyclopropyl-4-methylpentanoyl]-L-phenyl-alanine-N-methylamide**

The compound from Step 2 above (250 mg, 0.486 mmol) was taken up in ethanol (20 mL) and exposed to a hydrogen atmosphere in the presence of 10% Pd on charcoal (100 mg) for 2 h. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol, and the combined ethanol solution was evaporated to dryness to obtain the crude title compound as a white solid.

**Step 4: [2R-(1-Benzylhydroxycarbonyl)cyclopropyl-4-methyl-pentanoyl]-L-phenylalanine-N-methylamide**

The crude compound from Step 3 above was dissolved in a mixture of DMF (3 mL) and acetonitrile (12 mL). To this solution, O-benzylhydroxylamine hydrochloride (90 mg, 0.55 mmol), N-methylmorpholine (0.12 mL, 1.1 mmol), and O-benzotriazol-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU; 193 mg, 0.6 mmol) were added in this order. After stirring for 15 h at room temperature, the solvent was removed under reduced pressure and the residue was taken up in ethyl acetate. Sequential washing with 2N aqueous HCl, saturated aqueous sodium bicarbonate and water, followed by drying over Na<sub>2</sub>SO<sub>4</sub> and evaporation, left a residue which was purified by flash chromatography over silica (n-hexane and ethyl acetate gradient elution), thereby obtaining the title compound as a white solid (55 mg, 0.12 mmol; 25% over the two steps).

**Step 5: [2R-(1-Hydroxycarbonyl)cyclopropyl-4-methyl-pentanoyl]-L-phenylalanine-N-methylamide**

A solution of the compound from Step 4 above (55 mg, 0.12 mmol) in ethanol (5 mL) was exposed to a hydrogen atmosphere in the presence of 5% Pd on barium sulfate (30 mg) for 2.5 h. The catalyst was removed by filtration (Celite<sup>R</sup> filter aid), washed with additional ethanol, and the combined ethanol solution was evaporated to dryness. The residue was triturated with ethyl ether, and the resulting white powder was collected by filtration (20 mg; 0.053 mmol; 44%). IR (KBr),  $\nu_{\max}$  3380, 2958, 1654, and 1660 cm<sup>-1</sup>. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.18, 0.43, 0.54 and 0.98 (each m, 4 H, cyclopropyl -CH<sub>2</sub>-CH<sub>2</sub>), 0.77 and 0.78 (each d, J= 6.8 Hz, 6 H, 2 x Me), 1.16 and 1.68 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.33 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 2.09 (dd, J= 4.7 and 10.3 Hz, 1 H, CH-iBu), 2.55 (d, J= 4.7 Hz, 3 H, CONHCH<sub>3</sub>),


94

2.75 and 2.94 (each m, 2 H, Phe methylene), 4.50 (m, 1 H, CONHCH), 7.1-7.3 (m, 5 H, Ph), 7.97 (q,  $J = 4.7$  Hz, 1 H, CONHCH<sub>3</sub>), 8.47 (d,  $J = 9.0$  Hz, 1 H, CONHCH), 8.62 (s, 1 H, CONHOH), and 10.97 ppm (s, 1 H, CONHOH).

5

**EXAMPLE 21**

**[2R-[(2S)-2-Carboxy-2-oxiranyl]-4-methylpentanoyl]-L-phenylalanine-N-methylamide**




[2R-[(2S)-2-Benzhydryloxycarbonyl-2-oxiranyl]-4-methyl-  
 10 pentanoyl]-L-phenylalanine-N-methylamide (25 mg, 0.047 mmol),  
 prepared as described in Example 18, Step 1, was dissolved in  
 ethanol (5 mL) and exposed to a hydrogen atmosphere in the  
 presence of 10% Pd on charcoal (10 mg) for 2 h. The catalyst  
 was removed by filtration (Celite<sup>R</sup> filter aid), washed with  
 15 additional ethanol, and the combined ethanol solution was  
 evaporated to dryness. The residue was triturated with ethyl  
 ether, and the resulting white solid was collected by  
 filtration (15 mg, 0.041 mmol; 87%). <sup>1</sup>H NMR (400 MHz, DMSO-  
 d<sub>6</sub>): 0.67 and 0.70 (each d,  $J = 6.4$  Hz, 6 H, 2 x Me), 0.95-1.30  
 20 (m, 3 H, CH<sub>2</sub>CHMe<sub>2</sub>), 2.21 and 2.55 (each d,  $J = 6.0$  Hz, 2 H,  
 oxirane methylene), 2.51 (d,  $J = 4.7$  Hz, 3 H, CONHCH<sub>3</sub>), 2.74  
 (dd,  $J = 13.6$  and  $10.0$  Hz, 1 H, CH-CHH-Ph), 2.82 (m, 1 H, CH-  
 iBu), 2.96 (dd,  $J = 13.6$  and  $4.7$  Hz, 1 H, CH-CHH-Ph), 4.30 (m,  
 1 H, CONHCH), 7.1-7.3 (m, 5 H, Ph), 7.90 (m, 1 H, CONHCH<sub>3</sub>),  
 25 and 8.60 ppm (m, 1 H, CONHCH).

**EXAMPLE 22**



95

**1-[(2R,3S)-3-Benzoyloxymethyl-3-hydroxy-4-hydroxyamino-2-isobutylsuccinyl]morpholine**



**Step 1: 1-[(2R,3S)-4-Benzoyloxy-3-benzoyloxymethyl-3-hydroxy-2-isobutylsuccinyl]morpholine**

(2R,3S)-(4-Benzoyloxy-3-benzoyloxymethyl-3-hydroxy-2-isobutyl)-succinic acid (1.15 g, 2.77 mmol), obtained as described in Preparation 5, was dissolved in a mixture of dry DMF (8 mL) and dichloromethane (12 mL). To this solution, N-methyl-morpholine (0.39 mL), HOBT (1-hydroxybenzotriazole hydrate; 478 mg), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (670 mg) were added at 0°C. After 75 min, morpholine (0.15 mL) was added, and the reaction mixture was left for 20 hr at room temperature. The solution was taken up in ethyl acetate, and sequentially washed with 2% aqueous HCl, saturated aqueous sodium bicarbonate, and brine. After drying over Na<sub>2</sub>SO<sub>4</sub>, the solvent was removed and the residue was purified by flash-chromatography, thereby obtaining the title product as an oil (960 mg, 1.99 mmol; 72%).

**Step 2: 1-[(2R,3S)-3-Benzoyloxymethyl-1,4-dihydroxy-2-isobutylsuccinyl]morpholine**

A solution of the compound from Step 1 above (957 mg; 1.98 mmol) in ethanol (35 mL) was exposed to a hydrogen atmosphere in the presence of 5% Pd on charcoal (450 mg) for 2 hr. The catalyst was filtered off, washed with ethanol, and the

combined solution was evaporated to dryness. The residue was taken up in ethyl ether and concentrated to obtain the title product as a white powder (750 mg, 96%).

**Step 3: 1-[(2R,3S)-4-Benzoyloxyamino-3-benzoyloxymethyl-3-hydroxy-2-isobutylsuccinyl]morpholine**

A solution in acetonitrile (30 mL) of the compound from Step 2 above (750 mg, 1.91 mmol) was allowed to react with O-benzylhydroxylamine hydrochloride (335 mg, 2.01 mmol), N-methylmorpholine (0.46 mL, 4.2 mmol), and O-benzotriazolyl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU; 735 mg, 2.3 mmol). After 16 hr at room temperature, the solvent was removed under reduced pressure and the residue was taken up in ethyl acetate. Sequential washing with 2N aqueous HCl, saturated aqueous sodium bicarbonate and water, followed by drying over Na<sub>2</sub>SO<sub>4</sub> and evaporation, left a residue (800 mg) which was purified by flash chromatography over silica (n-hexane and ethyl acetate gradient elution), thereby obtaining the pure title product (504 mg, 1.01 mmol; 53%).

**Step 4: 1-[(2R,3S)-3-Benzoyloxymethyl-3-hydroxy-4-hydroxyamino-2-isobutylsuccinyl] morpholine**


A solution of the compound from Step 3 above (500 mg; 1 mmol) in ethanol (40 mL) was exposed to a hydrogen atmosphere in the presence of 5% Pd on charcoal (167 mg) for 3 hr. The catalyst was filtered off, washed with ethanol, and the combined solution was filtered through a Millipore<sup>R</sup> membrane. After evaporation to dryness, the residue was taken up in ethyl ether and concentrated to obtain the title product as a white solid (340 mg, 0.83 mmol; 83%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.81 and 0.82 (each d, J= 6.0 Hz, 6 H, 2 x Me), 1.22 and 1.78 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.30 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 3.30 (obscured by water, 1 H, CH-iBu), 3.40-3.60 (m, 8 H,

97

morpholino), 4.29 and 4.33 (each d,  $J = 10.7$  Hz,  $\text{CH}_2\text{OCOPh}$ ), 5.73 (s, 1 H, tertiary OH), 7.51 (m, 2 H, meta-benzoyl protons), 7.65 (m, 1 H, para-benzoyl proton), 7.92 (m, 2 H, ortho-benzoyl protons), 8.85 (s, 1 H,  $\text{NHOH}$ ), and 10.65 ppm (s, 1 H,  $\text{NHOH}$ ).

**EXAMPLE 23**


**1-[(2R,3S)-3-Acetoxymethyl-3-hydroxy-4-hydroxyamino-2-isobutylsuccinyl]piperidine**



The title compound was obtained starting from (2R,3S)-(3-acetoxymethyl-4-benzyloxy-3-hydroxy-2-isobutyl) succinic acid (see Preparation 4), by following the procedure described in Example 22 above but replacing morpholine with piperidine.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ ): 0.77 and 0.78 (each d,  $J = 6.8$  Hz, 6 H, 2 x Me), 1.05 and 1.65 (each m, 2 H,  $\text{CH}_2\text{CHMe}_2$ ), 1.23 (m, 1 H,  $\text{CH}_2\text{CHMe}_2$ ), 1.3-1.6 (m, 6 H, piperidine 3,4,5-methylene protons), 1.92 (s, 3 H,  $\text{COCH}_3$ ), 3.21 (dd,  $J = 3.0$  and 11.1 Hz, 1 H,  $\text{CH}-i\text{Bu}$ ), 3.50 (m, 4 H, piperidine 2,6-methylene protons), 3.92 and 4.00 (each d,  $J = 11.1$  Hz, 2 H,  $\text{CH}_2\text{OCO}$ ), 5.65 (s, 1 H, tertiary OH), 8.75 (s, 1 H,  $\text{NHOH}$ ), and 10.48 ppm (s, 1 H,  $\text{NHOH}$ ).

**EXAMPLE 24**

**1-[(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutyl-succinyl]piperidine**

**ROUTE A**

**Steps 1-3: 1-[(2R,3S)-(3-Acetoxymethyl-4-benzyloxyamino-3-hydroxy-2-isobutyl)succinyl]piperidine**

The title compound was obtained starting from (2R,3S)-(3-acetoxymethyl-4-benzyloxy-3-hydroxy-2-isobutyl)succinic acid (see Preparation 4), and following the first 3 steps of the preparation of Example 23.

**Step 4: 1-[(2R,3S)-(4-Benzyloxyamino-3-hydroxy-3-hydroxymethyl-2-isobutyl)succinyl] piperidine**

A solution of the compound from Step 3 above (600 mg) in methanol (40 mL) was allowed to react with sodium methoxide (112 mg) for 2 hr at room temperature. After quenching with acetic acid (0.16 mL), the reaction mixture was partitioned between ethyl acetate and water. Drying of the organic extract with sodium sulfate and evaporation left the crude title compound as a white solid (540 mg).

**Step 5: 1-[(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-isobutyl-succinyl]piperidine**

A solution of the compound from Step 4 above (540 mg) in ethanol (40 mL) was exposed to a hydrogen atmosphere in the presence of 5% Pd/C (175 mg) for 90 min at room temperature. Filtration of the catalyst and evaporation to dryness afforded the title compound as a gummy solid (350 mg). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.76 and 0.77 (each d, J= 6.4 Hz, 6 H, 2 x Me), 1.02

99


and 1.65 (each m, 2 H,  $\text{CH}_2\text{CHMe}_2$ ), 1.20 (m, 1 H,  $\text{CH}_2\text{CHMe}_2$ ), 1.3-1.6 (m, 6 H, piperidine 3,4,5-methylene protons), 3.20 (m, 2 H,  $\text{CH-iBu}$  and  $\text{CH(H)OH}$ ), 3.3-3.5 (m, 5 H, piperidine 2,6-methylene protons and  $\text{CH(H)OH}$ ), 4.63 (broad s, 1 H,  $\text{CH}_2\text{OH}$ ),  
5 5.38 (s, 1 H, tertiary OH), 8.63 (broad s, 1 H,  $\text{NHOH}$ ), and 10.20 ppm (broad s, 1 H,  $\text{NHOH}$ ).

**ROUTE B**

A solution in dry DMF of (2R,3S)-(3-benzyloxyaminocarbonyl-3-hydroxy-2-isobutyl)butyrolactone (400 mg, obtained as  
10 described in Preparation 8) was allowed to react with piperidine (0.5 mL) for 2 days. The reaction mixture was partitioned between ethyl acetate and 4% aqueous HCl. After washing with saturated sodium hydrogen carbonate and drying over sodium sulfate, the solvent was removed in vacuo and the  
15 residue purified by flash-chromatography over silica, thereby obtaining 1-[(2R,3S)-(4-benzyloxyamino-3-hydroxy-3-hydroxymethyl-2-isobutyl)succinyl]piperidine, identical with the material obtained by Route A, Step 4. Hydrogenolysis of this compound, as described in Step 5, Route A, afforded the title  
20 compound.

**EXAMPLE 25**

**1-[(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(trimethylacetoxymethylsuccinyl)morpholine**



**Step 1: 1-[(2R,3S)-4-Benzylloxy-3-hydroxy-2-isobutyl-3-(trimethylacetoxymethylsuccinyl)morpholine**

N-Methylmorpholine (0.382 mL) and HOBt (465 mg) were added to a solution in dichloromethane (12 mL) of (2R,3S)-[4-benzylloxy-3-hydroxy-2-isobutyl-3-(trimethylacetoxymethyl)succinic acid (1.13 g), obtained as described in Example 13, Step 1. After cooling to 0 °C, N, N-dimethylaminopropyl-N'-ethylcarbodiimide hydrochloride (660 mg) was added, followed after 75 min at 0 °C by morpholine (0.5 mL). The reaction mixture was left standing for 2 days at room temperature. The solution was taken up in ethyl acetate, and sequentially washed with 2% aqueous HCl, saturated aqueous sodium bicarbonate, and brine. After drying over Na<sub>2</sub>SO<sub>4</sub>, the solvent was removed and the residue was purified by flash-chromatography, thereby obtaining the title product as a yellowish oil (1.28 g).

**Step 2: 1-[(2R,3S)-3,4-Dihydroxy-2-isobutyl-3-(trimethylacetoxymethylsuccinyl)morpholine**

A solution of the compound from Step 1 above (1.25 g) in ethanol (40 mL) was exposed to a hydrogen atmosphere in the presence of 5% Pd on charcoal (0.5 g) for 2 hr. The catalyst was filtered off, washed with ethanol, and the combined solution was evaporated to dryness. The residue was taken up in ethyl ether and concentrated to obtain the title product as a gummy solid (970 mg).

**Step 3: 1-[(2R,3S)-4-Benzylloxyamino-3-hydroxy-2-isobutyl-3-(trimethyl)acetoxymethylsuccinyl)morpholine**

A solution in acetonitrile (30 mL) of the compound from Step 2 above (950 mg) was allowed to react with O-benzylhydroxylamine hydrochloride (456 mg), N-methylmorpholine (0.62 mL), and TBTU (735 mg). After 40 hr at room temperature, the solution was concentrated under reduced pressure to a small volume and

101

partitioned between ethyl acetate and water. Sequential washing with 2N aqueous HCl, saturated aqueous sodium bicarbonate and water, followed by drying over Na<sub>2</sub>SO<sub>4</sub> and evaporation, left a residue (800 mg) which was purified by  
5 flash chromatography over silica (n-hexane and ethyl acetate gradient elution), thereby obtaining the pure title product (580 mg) as a white powder.

**Step 4: 1-[(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(trimethylacetoxymethylsuccinyl)morpholine**

10 A solution of the compound from Step 3 above (580 mg) in ethanol (40 mL) was exposed to a hydrogen atmosphere in the presence of 5% Pd on charcoal (275 mg) for 2 hr. The catalyst was filtered off, washed with ethanol, and the combined solution was evaporated to dryness. The residue was triturated  
15 in a small volume of ethyl ether to obtain the title product as a white solid (460 mg). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.78 and 0.80 (each d, J= 6.5 Hz, 6 H, 2 x Me), 1.09 (s, 9 H, t-Bu), 1.0-1.8 (m, 3 H, CH<sub>2</sub>CHMe<sub>2</sub>), 3.3 (obscured by water, 1 H, CH-iBu), 3.57 (m, 8 H, morpholino), 4.00 and 4.08 (each d, J= 11.0 Hz, CH<sub>2</sub>OCO), 5.43 (s, 1 H, tertiary OH), 8.75 (s, 1 H, NHOH), and 10.43 ppm (s, 1 H, NHOH).

**EXAMPLE 26**

Similarly, the following compounds may be prepared:

#26.1: 1-[2(R)-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]piperidine;

#26.2: 1-[2(R)-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-mercaptomethylsuccinyl]piperidine;

#26.3: 1-[(2R,3S)-3-Hydroxy-4-hydroxyamino-3-hydroxymethyl-2-octylsuccinyl]pyridazine;

- #26.4: 1-[(2R,3S)-2-(3-(4-Biphenyl)propyl)-3-hydroxy-4-hydroxy-amino-3-hydroxymethylsuccinyl]-6-(N-methylcarbamoyl)-hexahydropyridazine;
- #26.5: 1-[(2R,3S)-3-Hydroxy-4-hydroxyamino-2-isobutyl-3-(1-methyl-1,2,3,4-tetrazol-5-yl)thiomethylsuccinyl]piperidine;
- #26.6: 1-[(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-(4-hydroxyphenyl)thiomethylsuccinyl]piperidine;
- #26.7: 1-[(2R,3S)-2-Cyclopentylmethyl-3-(4-fluorobenzene-sulfonyl)methyl-3-hydroxy-4-hydroxyaminosuccinyl]piperidine;
- #26.8: 1-[(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxyamino-3-(4-methoxybenzenesulfonyl)methylsuccinyl]-6-(N-methyl-carbamoyl)hexahydropyridazine;
- #26.9: 1-[(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-2-cyclopentylmethyl-3-hydroxy-4-hydroxyaminosuccinyl]piperidine;
- #26.10: 1-[(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-2-cyclopentylmethyl-3-hydroxy-4-hydroxyaminosuccinyl]-6-(N-methyl-carbamoyl)hexahydropyridazine;
- #26.11: N-[(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-2-cyclopentylmethyl-3-hydroxy-4-hydroxyaminosuccinyl]-2-cyclohexylethylamine;
- #26.12: (R)-N-[(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-2-cyclopentylmethyl-3-hydroxy-4-hydroxyaminosuccinyl]-1-cyclohexylethylamine;
- #26.13: (R)-N-[(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-2-cyclopentylmethyl-3-hydroxy-4-hydroxyaminosuccinyl]-1-phenyl-ethylamine;
- #26.14: (R)-N-[(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-2-cyclopentylmethyl-3-hydroxy-4-hydroxyaminosuccinyl]-1-benzyl-ethylamine;



- #26.15: (S)-N-[(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-2-cyclopentylmethyl-3-hydroxy-4-hydroxyaminosuccinyl]-1-benzyl-2-hydroxyethylamine;
- #26.16: (S)-N-[(2R,3S)-2-(3-(4-Chlorophenyl)propyl)-3-(N-cyclohexylaminocarbonyl)oxymethyl-3-hydroxy-4-hydroxyamino-succinyl]-1-benzyl-2-hydroxyethylamine;
- #26.17: (S)-N-[(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-2-cyclopentylmethyl-3-hydroxy-4-hydroxyaminosuccinyl]-1-(1H-imidazol-5-yl)methyl-2-hydroxyethylamine;
- #26.18: (S)-N-[(2R,3S)-2-(3-(4-Biphenyl)propyl)-3-hydroxy-4-hydroxyamino-3-hydroxymethylsuccinyl]-1-(1H-imidazol-5-yl)-methyl-2-hydroxyethylamine;
- #26.19: (S)-N-[(2R,3S)-2-(3-(4-Biphenyl)propyl)-3-(N-cyclohexylaminocarbonyl)oxymethyl-3-hydroxy-4-hydroxyaminosuccinyl]-1-(1H-imidazol-5-yl)methyl-2-hydroxyethylamine;
- #26.20: (S)-N-[(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxy-amino-3-hydroxymethylsuccinyl]-(3,4-dihydro) carbostyryl-3-amine;
- #26.21: (S)-N-[(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxy-amino-3-hydroxymethylsuccinyl]-(N-methyl-3,4-dihydro)-carbostyryl-3-amine;
- #26.22: (S)-N-[(2R,3S)-2-Cyclopentylmethyl-3-hydroxy-4-hydroxy-amino-3-hydroxymethylsuccinyl]-(6,7-dioxymethylene-N-ethyl-3,4-dihydro) carbostyryl-3-amine;
- #26.23: (S)-N-[(2R,3S)-3-(N-Cyclohexylaminocarbonyl)oxymethyl-2-cyclopentylmethyl-3-hydroxy-4-hydroxyaminosuccinyl]-(6,7-dioxymethylene-N-ethyl-3,4-dihydro) carbostyryl-3-amine.

#### Example 27

- Tablets containing the following ingredients may be produced in a conventional manner:

104

	Ingredient	Per Tablet
	Compound of formula (I)	25.0 mg
	Lactose	125.0 mg
	Maize starch	75.0 mg
5	Talc	4.0 mg
	Magnesium stearate	1.0 mg
	Total weight	230.0 mg


**Example 28**

Capsules containing the following ingredients may be produced  
 10 in a conventional manner:

	Ingredient	Per capsule
	Compound of formula (I)	50.0 mg
	Lactose	165.0 mg
	Maize starch	20.0 mg
15	Talc	5.0 mg
	Capsule fill weight	240.0 mg

**PREPARATION 1**

*Preparation of benzyl (3R)-(2-benzyloxycarbonyl-3-carboxy-5-methyl)hexanoate by optical resolution.*



20

**Step 1: 2-Bromo-4-methylpentanoic acid**

D,L-Leucine (50 g, 0.38 mol) and potassium bromide (158.7 g, 1.33 mol) were dissolved in aqueous sulfuric acid (from 75 mL of H<sub>2</sub>SO<sub>4</sub> and 250 mL of water), and sodium nitrite (34.8 g, 0.5  
 25 mol) was added portionwise while keeping the temperature below -1 °C. The mixture was kept at 0°C for 1 h, then dichloromethane was added under stirring. The dichloromethane

105

layer was collected and the aqueous phase was extracted twice with further portions of dichloromethane. The combined dichloromethane extracts were dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated to give the title compound as a pale yellow oil (63.4 g, 0.32 mol; 85.5%).

**Step 2: *tert*-Butyl 2-bromo-4-methylpentanoate**

The product from Step 1 above (63.3 g, 0.32 mol) was dissolved in dichloromethane (250 mL), concentrated sulfuric acid (2 mL) was added, and cooled to  $-40^\circ\text{C}$  in a pressure bottle. Isobutene was condensed into the solution to roughly double the volume, and the mixture was then allowed to warm to room temperature overnight. Following evaporation to half of the volume under reduced pressure, the solution was washed with 10% aqueous sodium bicarbonate, dried over  $\text{Na}_2\text{SO}_4$ , and thoroughly evaporated under reduced pressure at room temperature to leave the title compound as a yellow oil (79.9 g, 0.314 mol; 98%).

**Step 3: *Benzyl* (2-benzyloxycarbonyl-3-*tert*-butoxycarbonyl-5-methyl)hexanoate**

Potassium *tert*-butoxide (26.9 g, 0.24 mol) was added portionwise under stirring to a cooled ( $-5^\circ\text{C}$ ) solution of dibenzyl malonate (59.3 mL, 0.24 mol) in dry DMF (100 mL). After further 10 min stirring between  $-5^\circ\text{C}$  and  $0^\circ\text{C}$ , a solution of *tert*-butyl 2-bromo-4-methylpentanoate (60.7 g, 0.24 mol) from Step 2 above in dry DMF (100 mL) was added dropwise over 1 h while keeping the temperature at  $0^\circ\text{C}$ . After the addition was complete, the reaction mixture was kept for 4 days in a cold room ( $4^\circ\text{C}$ ), then partitioned between ethyl acetate and saturated aqueous ammonium chloride. The aqueous layer was extracted twice with further ethyl acetate before being discarded. The combined organic extracts were dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated, to left a residue which was purified by flash

chromatography over silica (95:5 n-hexane and ethyl acetate). The title compound was obtained as a colourless oil (79.9 g, 0.175 mol; 73.1%). <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>): 0.83 and 0.84 (each d, J= 6.5 Hz, 6 H, 2 x Me), 1.0-1.6 (m, 3 H, CH<sub>2</sub>CHMe<sub>2</sub>),  
5 1.39 (s, 9 H, tert-Bu), 3.06 (m, 1 H, CH-iBu), 3.74 (d, J= 10.1 Hz, 1 H, CH(CO<sub>2</sub>Bn)<sub>2</sub>), 5.12 (m, 4 H, 2 x CO<sub>2</sub>CH<sub>2</sub>Ph), and 7.30 ppm (m, 10 H, 2 x Ph).

**Step 4: Benzyl (2-benzyloxycarbonyl-3-carboxy-5-methyl)-hexanoate**

10 The compound from Step 3 above (72 g, 0.16 mol) was added to a solution of trifluoroacetic acid (100 mL) and water (5 mL) and allowed to stand at 4°C overnight. After evaporation under reduced pressure, the residue was partitioned between dichloromethane (250 mL) and brine (250 mL). The organic layer  
15 was dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to give the crude title product as a colourless oil (63.7 g, 0.16 mol; quantitative yield).

**Step 5: Optical resolution of Benzyl (3R)-(2-benzyloxycarbonyl-3-carboxy-5-methyl)hexanoate**


20 The racemic crude compound from Step 4 above (63.7 g, 0.16 mol) was taken up in diisopropyl ether (300 mL) and the mixture heated to reflux temperature until a homogeneous solution formed, which while hot was treated dropwise with S(-)-1-phenylethylamine (10.3 mL, 0.08 mol). The mixture was then  
25 allowed to cool to room temperature; filtration gave the phenylethylammonium salt of the title (3R) acid (36.4 g, 0.07 mol). This salt was added under stirring to a mixture of ethyl ether (250 mL) and 2N aqueous HCl (35 mL, 0.07 mol). The aqueous layer was extracted with further portions of ethyl  
30 ether, and the combined ethereal layers were dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to give the title compound as a colourless oil

107

which crystallized on standing (27.9 g, 0.07 mol, 87.5%).  $^1\text{H}$  NMR (200 MHz,  $\text{DMSO-d}_6$ ): 0.76 (d,  $J = 6.4$  Hz, 6 H, 2 x Me), 1.0-1.6 (m, 3 H,  $\text{CH}_2\text{CHMe}_2$ ), 2.92 (m, 1 H,  $\text{CH-iBu}$ ), 3.70 (d,  $J = 10.1$  Hz, 1 H,  $\text{CH}(\text{CO}_2\text{Bn})_2$ ), 5.11 (s, 2 H,  $\text{CO}_2\text{CH}_2\text{Ph}$ ), 5.12 and 5.17 (each d,  $J = 12.3$  Hz, 2 H,  $\text{CO}_2\text{CH}_2\text{Ph}$ ), 7.32 (m, 10 H, 2 x Ph), and 12.59 ppm (broad s, 1 H,  $\text{CO}_2\text{H}$ ).

## PREPARATION 2

***tert*-Butyl (2R,3S)-(3,4-dihydroxy-3-hydroxymethyl-2-isobutyl)-succinate**



10

**Step 1: *tert*-Butyl (2R)-(4-hydroxy-2-isobutyl-3-methylene)-succinate**

A solution in ethanol (250 mL) of benzyl (3R)-(2-benzyloxycarbonyl-3-carboxy-5-methyl)hexanoate (60 g; 0.13 mol), obtained as described in Preparation 1, was exposed to a hydrogen atmosphere in the presence of 5% Pd on charcoal (20 g) overnight. The catalyst was filtered off, washed with ethanol, and piperidine (15 mL) and 37% aqueous formaldehyde (91 mL) were added to the combined ethanol solution. After standing at 4°C for 3 days, the reaction mixture was concentrated in vacuo, taken up with ethyl acetate, and washed with 4% aqueous HCl. After drying over  $\text{Na}_2\text{SO}_4$  and evaporation, the crude title product was obtained (32 g, 0.13 mol; quantitative yield).


**Step 2: *tert*-Butyl (2R,3S)-(3,4-dihydroxy-3-hydroxymethyl-2-isobutyl)succinate**

108

N-Methylmorpholine-N-oxide (18.5 g, 0.159 mol) was dissolved in a mixture of water (10 mL) and acetone (5 mL), and osmium tetroxide (2.5% in tert-butanol; 13 mL, 1.32 mmol) was added. After stirring for 15 min, a solution of the crude compound from Step 1 above (32 g, 0.13 mol) in tert-butanol (300 mL) was added. After stirring for 18 h at room temperature, the solution was concentrated and the residue was taken up in ethyl acetate, washed with 4% aqueous HCl, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness to leave a brownish oil. Addition of ethyl ether and stirring caused the separation of the pure title product, which was collected as a white powder by filtration (8.6 g). Removal of the solvent from the mother liquors gave a further crop of the title compound as a syrup (23.25; total 31.85 g, 0.116 mol; 87.9%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.79 and 0.82 (each d, J= 6.4 Hz, 6 H, 2 x Me), 0.95 and 1.77 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.36 (s, 9 H, t-Bu), 1.36 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 2.57 (dd, J= 2.9 and 12.0 Hz, 1 H, CH-iBu), 3.39 and 3.65 ppm (each d, J= 10.8 Hz, 2 H, CH<sub>2</sub>OH).

## PREPARATION 3

**tert-Butyl (2R,3S)-(4-benzoyloxy-3-hydroxy-3-hydroxymethyl-2-isobutyl)succinate**




A solution) in dry DMF (210 mL) of tert-butyl (2R,3S)-(3,4-dihydroxy-3-hydroxymethyl-2-isobutyl)succinate (23.2 g, 96 mmol), obtained as described in Preparation 2, was treated at 0°C with benzyl bromide (12.5 mL, 106 mmol) and triethylamine

109

(14.7 mL, 106 mmol). After 15 min at 0°C, the cooling bath was removed and the mixture was stirred for 20 hr. Partition between water and ethyl acetate, followed by separation of the organic layer, sequential washing with 4% aqueous HCl, saturated aqueous sodium bicarbonate and water, drying (Na<sub>2</sub>SO<sub>4</sub>) and evaporation, left a residue, which was fractionated by flash chromatography over silica (4:1 hexane - ethyl acetate). The title product was obtained as the slower-eluting compound (13.5 g, 37 mmol; 38.5%). A second crop, consisting of a mixture of the same compound with the 2R,3R isomer, was also obtained (7.54 g, 20.6 mmol; 21.5%; total yield 60%). IR (CHCl<sub>3</sub> film),  $\nu_{\max}$  2959, 2872, and 1738 cm<sup>-1</sup>. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.67 and 0.75 (each d, J= 6.6 Hz, 6 H, 2 x Me), 0.81 and 1.74 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.31 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.35 (s, 9 H, t-Bu), 2.58 (dd, J= 2.6 and 11.9 Hz, 1 H, CH-iBu), 3.45 (dd, J= 4.8 and 11.0 Hz, 1 H, CH(H)OH), 3.70 (dd, J= 6.8 and 11.0 Hz, 1 H, CH(H)OH), 4.90 (dd, J= 4.8 and 6.8 Hz, 1 H, primary OH), 5.01 (s, 1 H, tertiary OH), 5.11 and 5.16 (each d, J= 12.5 Hz, 2 H, COOCH<sub>2</sub>Ph), and 7.36 ppm (m, 5 H, Ph).

**PREPARATION 4**

**(2R,3S)-(3-Acetoxymethyl-4-benzyloxy-3-hydroxy-2-isobutyl)-succinic acid**



110

**Step 1: tert-Butyl (2R,3S)-(3-acetoxymethyl-4-benzyloxy-3-hydroxy-2-isobutyl)succinate**


tert-Butyl (2R,3S)-(4-benzyloxy-3-hydroxy-3-hydroxymethyl-2-isobutyl)succinate (2.1 g, 5.75 mmol), obtained as described  
5 in Preparation 3, was dissolved in dichloromethane (80 mL). Pyridine (0.92 mL, 11.5 mmol) and then acetyl chloride (0.81 mL, 11.5 mmol) was added dropwise at 0°C. The cooling bath was removed and the solution was left for 5 hr under stirring. Sequential washing with 4% aqueous HCl, saturated aqueous  
10 sodium bicarbonate and water, drying (Na<sub>2</sub>SO<sub>4</sub>) and evaporation, left the title product (essentially pure by HPLC; 2.16 g, 5.3 mmol; 92.2%).

**Step 2: (2R,3S)-(3-Acetoxymethyl-4-benzyloxy-3-hydroxy-2-isobutyl)succinic acid**

15 Trifluoroacetic acid (2.3 mL) was added to a cooled (0°C) solution of the compound from Step 1 above (2.16 g, 5.3 mmol) in dichloromethane (5 mL). The cooling bath was removed, and the mixture was stirred overnight. The solution was repeatedly evaporated in vacuo after repeated addition of ethyl ether, thereby obtaining the crude title compound as an oil (1.87 g,  
20 5.3 mmol; quantitative yield).

**PREPARATION 5**

**(2R,3S)-(4-Benzyloxy-3-benzoyloxymethyl-3-hydroxy-2-isobutyl)-succinic acid**





111

**Step 1: *tert*-Butyl (2*R*,3*S*)-(4-benzyloxy-3-benzoyloxymethyl-3-hydroxy-2-isobutyl)succinate**

A solution in dry dichloromethane (30 mL) of *tert*-butyl (2*R*,3*S*)-(4-benzyloxy-3-hydroxy-3-hydroxymethyl-2-isobutyl)-  
5 succinate (1.83 g, 5 mmol), obtained as described in Preparation 3, was sequentially treated at 0°C with benzoyl chloride (0.7 mL, 6 mmol) and triethylamine (0.84 mL, 6 mmol). The solution was left aside for 20 hr at room temperature. Aqueous work-up (washing with sodium bicarbonate and diluted  
10 HCl) and flash chromatography afforded the title compound as a white powder (1.4 g, 3 mmol; 60%).


**Step 2: (2*R*,3*S*)-(4-Benzyloxy-3-benzoyloxymethyl-3-hydroxy-2-isobutyl)succinic acid**

The compound from Step 1 above (1.4 g, 3 mmol) was dissolved  
15 in dichloromethane (10 mL) and treated with trifluoroacetic acid (4 mL). After 2 hr at room temperature, the reaction mixture was evaporated to dryness, and the residue was triturated with isopropyl ether and filtered, thereby obtaining the title compound as a white powder (550 mg). A  
20 second crop (600 mg) was recovered from the mother liquors by evaporation and flash-chromatography over silica (*n*-hexane and ethyl acetate as eluants). Total yield 1.15 g (2.8 mmol; 92%).  
<sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>): 0.77 and 0.84 (each d, J= 6.4 Hz, 6 H, 2 x Me), 0.92 and 1.90 (each m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.51 (m, 1  
25 H, CH<sub>2</sub>CHMe<sub>2</sub>), 3.03 (dd, J= 3.2 and 11.9 Hz, 1 H, CH-*i*Bu), 4.58 (s, 2 H, CH<sub>2</sub>OCOPh), 5.23 and 5.30 (each d, J= 11.7 Hz, 2 H, COOCH<sub>2</sub>Ph), and 7.2-7.9 ppm (m, 10 H, Ph).

**PREPARATION 6**

**(2*R*,3*S*)-(2-Benzyl-3-carboxy-3-hydroxy)butyrolactone**


112



Trifluoroacetic acid (2 mL) was added to a solution of tert-butyl (2R,3S)-(2-benzyl-3,4-dihydroxy-3-hydroxymethyl) succinate (300 mg; 1.02 mmol) in dichloromethane (6 mL). After 3 hr at room temperature, the mixture was concentrated in vacuo to leave a residue. Trituration with isopropyl ether and filtration gave the title product as a white powder (180 mg, 0.76 mmol; 75%). <sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>): 2.82 (m, 2 H, CH<sub>2</sub>Ph), 3.30 (m, 1 H, CHCH<sub>2</sub>Ph), 4.11 and 4.28 (each d, J= 9.4 Hz, 2 H, lactone methylene), 6.10 (broad s, 1 H, OH), 7.17 (m, 5 H, Ph), and 13.00 ppm (broad s, 1 H, COOH).

#### PREPARATION 7

(2R,3S)-(2-Benzyl-3-benzyloxyaminocarbonyl-3-hydroxy)-butyrolactone




By following a procedure analogous to that described in Example 2, Step 1, a solution in acetonitrile (10 mL) of (2R,3S)-(2-benzyl-3-carboxy-3-hydroxy)butyrolactone (110 mg, 0.46 mmol), obtained as described in Preparation 6, was allowed to react with O-benzylhydroxylamine hydrochloride (88

113

mg), N-methylmorpholine (0.12 mL), and O-benzotriazolyl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU; 190 mg). After 2 days at room temperature, work-up and flash chromatography over silica afforded the title product as a white powder (85 mg, 0.25 mmol; 54%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 2.77 (dd, J= 9.0 and 14.5 Hz, 1 H, CH(H)Ph), 2.86 (dd, J= 4.7 and 14.5 Hz, 1 H, CH(H)Ph), 3.38 (dd, J= 4.7 and 9.0 Hz, 1 H, CHCH<sub>2</sub>Ph), 4.18 and 4.31 (each d, J= 9.8 Hz, 2 H, lactone methylene), 4.15 and 4.23 (each d, J= 10.2 Hz, 2 H, OCH<sub>2</sub>Ph), 6.53 (s, 1 H, OH), 7.0-7.3 (m, 10 H, 2 x Ph), and 11.31 ppm (s, 1 H, CONHO).

#### PREPARATION 8

**(2R,3S)-(3-Benzylxyaminocarbonyl-3-hydroxy-2-isobutyl)-butyrolactone**




Starting from tert-butyl (2R,3S)-(3,4-dihydroxy-3-hydroxymethyl-2-isobutyl)succinate (0.6 g), obtained as described in Preparation 2, and using identical procedures to those described in Preparation 6 and Preparation 7, in this order, the title compound was obtained as a white powder (0.3 g).

#### PREPARATION 9

**(2R,3S)-(3-Acetoxymethyl-3-benzylxycarbonyl-2-isobutyl)-propiolactone**

114



A solution in dichloromethane (10 mL) of (2R,3S)-(3-acetoxymethyl-4-benzyloxy-3-hydroxy-2-isobutyl)succinic acid, (210 mg), obtained as described in Preparation 4, was allowed to react with N, N-dimethylaminopropyl-N'-ethylcarbodiimide hydrochloride (120 mg) for 15 hr at room temperature. The reaction mixture was washed with water and evaporated to leave the crude title product as a waxy solid in quantitative yield.


10 An analytical sample was obtained by flash chromatography over silica. IR (CHCl<sub>3</sub> film),  $\nu_{\max}$  1831, 1772, and 1713 cm<sup>-1</sup>. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 0.74 and 0.78 (each d, J= 6.7 Hz, 6 H, 2 x Me), 1.26 (m, 2 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.56 (m, 1 H, CH<sub>2</sub>CHMe<sub>2</sub>), 1.95 (s, 3 H, COMe), 4.15 (dd, J= 6.3 and 9.0 Hz, 1 H, CH-iBu), 4.54 and 4.69 (each d, J= 12.8 Hz, 2 H, CH<sub>2</sub>OAc), 5.27 and 5.32 (each d, J= 12.1 Hz, 2 H, COOCH<sub>2</sub>Ph), and 7.38 ppm (m, 5 H, Ph).

15

## CLAIMS

1. A compound of formula (I)

5



wherein


W is -NHOH or -OH;

10  $R_1$  is hydroxymethyl or a hydroxymethyl derivative which is an ether, an ester, a carbonate or a carbamate; or


$R_1$  is mercaptomethyl or a mercaptomethyl derivative which is a sulfide, a sulfoxide, a sulfone or a thioester;

$R_2$  is hydroxy or protected hydroxy; or


15  $R_1$  and  $R_2$ , taken together with the carbon atom to which they are attached, represent an optionally substituted cyclopropane, oxirane, 1,3-dioxolane or 2-oxo-1,3-dioxolane ring of the following formulae (B1)-(B4):




(B1)



(B2)



(B3)




(B4)

20


$R_3$  is a group  $-A^1-X-(CH_2)_n-A$ , wherein A is  $C_1-C_{10}$  alkyl,  $C_2-C_{10}$  alkenyl,  $C_3-C_7$  cycloalkyl, aryl, or heterocyclyl, the said

116

alkyl, alkenyl, cycloalkyl, aryl and heterocyclyl groups being unsubstituted or substituted; n is either zero or an integer from 1 to 5; -X- is either a direct bond or a group -O-, -S-, -SO-, -SO<sub>2</sub>-, -SO<sub>2</sub>NH-, -CO-, -CONH-, -NHCO-, -OCONH-, NHCONH or -NHSO<sub>2</sub>-, and -A<sup>I</sup>- is C<sub>1</sub>-C<sub>10</sub> alkylene, C<sub>2</sub>-C<sub>6</sub> alkenylene, or phenylene;  
 5 R<sub>4</sub> is a group of formula (C):




10 wherein R<sub>6</sub> is hydrogen or the side chain of a natural or non-natural alpha-amino acid, and R<sub>7</sub> is amino or a group -NH-A, -NH-CH<sub>2</sub>-A or -NH-CH<sub>2</sub>CH<sub>2</sub>-A wherein A is as defined above; or  
 R<sub>4</sub> is a group A as defined above or a group -A<sup>I</sup>-X-A wherein A, -X- and -A<sup>I</sup>- are as defined above;  
 15 or R<sub>4</sub> is a group of formula (D):



wherein R<sub>8</sub> is methyl, ethyl, phenyl, pyridyl, isopropyl, sec-butyl, tert-butyl, adamantyl, cyclopentyl, cyclohexyl, cyclohexylmethyl, phenylmethyl, 3,4-(methylenedioxy)phenyl,  
 20 piperonyl, pyridylmethyl, thienylmethyl, 3-indolylmethyl, 3-(N-methyl)indolylmethyl, 1,1-diphenylmethyl, naphthyl, naphthylmethyl, quinolylmethyl, isoquinolylmethyl, thiazolyl, thiazolylmethyl, imidazolyl, imidazolylmethyl, or a

117


derivative thereof optionally substituted, or  $R_8$  is -  
 $C(CH_3)_2SCH_3$  or a sulfoxide or sulfone thereof,  $-C(CH_3)_2OCH_3$ , -  
 $CH_2CH_2CH_2-OCH_3$ , or  $-CH(CH_3)OH$  or a tert-butyl ether thereof;  $R_9$   
 5 is either hydrogen or a group selected from methyl, ethyl,  
 phenyl, 3,4-(methylenedioxy)phenyl, piperonyl, pyridyl,  
 benzimidazolyl and 4-tetrazolyl, which group is optionally  
 substituted; or  $R_8$  and  $R_9$ , taken together with the nitrogen  
 atom to which they are attached, constitute a  
 dihydrocarbostyryl ring (D'):



10

wherein the nitrogen atom and the phenyl ring may be  
 optionally substituted;  
 $R_5$  is hydrogen or methyl; or  
 $R_4$  and  $R_5$ , taken together with the nitrogen atom to which  
 15 they are attached, form an optionally substituted  
 azaheterocyclyl ring;  
 and the solvates, hydrates and pharmaceutically acceptable  
 salts thereof.

2. A compound according to claim 1 wherein the absolute  
 20 stereochemistry of the succinic amide carbon atoms is the one  
 of the formula (I'):



3. A compound according to claim 2 wherein:

W is -NHOH or -OH;

R<sub>1</sub> is hydroxymethyl or the ether, ester, carbonate and carbamate derivatives thereof, as defined in claim 1; or

5 R<sub>1</sub> is a mercaptomethyl or the sulfide, sulfone and thioester derivatives thereof, as defined in claim 1;

R<sub>2</sub> is hydroxy or a protected derivative thereof; or

R<sub>1</sub> and R<sub>2</sub>, taken together with the carbon atom to which they are attached, form a cyclopropane or oxirane ring, optionally  
10 substituted by phenyl or benzyl; and:


R<sub>3</sub> is -CH<sub>2</sub>-alkyl, -CH<sub>2</sub>-cycloalkyl, -(CH<sub>2</sub>)<sub>n</sub>-O-alkyl, -(CH<sub>2</sub>)<sub>n</sub>-O-cycloalkyl, -(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>n'</sub>-aryl, -(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>n'</sub>-heterocyclyl, -(CH<sub>2</sub>)<sub>n</sub>-S-alkyl, -(CH<sub>2</sub>)<sub>n</sub>-S-cycloalkyl, -(CH<sub>2</sub>)<sub>n</sub>-S-(CH<sub>2</sub>)<sub>n'</sub>-aryl, -(CH<sub>2</sub>)<sub>n</sub>-S-(CH<sub>2</sub>)<sub>n'</sub>-heterocyclyl, -(CH<sub>2</sub>)<sub>n</sub>-S(O)-alkyl, -(CH<sub>2</sub>)<sub>n</sub>-S(O)-cycloalkyl, -(CH<sub>2</sub>)<sub>n</sub>-SO<sub>2</sub>-alkyl, -(CH<sub>2</sub>)<sub>n</sub>-SO<sub>2</sub>-cycloalkyl, -(CH<sub>2</sub>)<sub>n</sub>-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>n'</sub>-aryl, -(CH<sub>2</sub>)<sub>n</sub>-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>n'</sub>-heterocyclyl, -(CH<sub>2</sub>)<sub>n</sub>-SO<sub>2</sub>NH-alkyl, -(CH<sub>2</sub>)<sub>n</sub>-SO<sub>2</sub>NH-cycloalkyl, -(CH<sub>2</sub>)<sub>n</sub>-SO<sub>2</sub>NH-(CH<sub>2</sub>)<sub>n'</sub>-aryl, -(CH<sub>2</sub>)<sub>n</sub>-SO<sub>2</sub>NH-(CH<sub>2</sub>)<sub>n'</sub>-heterocyclyl, -(CH<sub>2</sub>)<sub>n</sub>-CO-alkyl, -(CH<sub>2</sub>)<sub>n</sub>-CO-cycloalkyl, -(CH<sub>2</sub>)<sub>n</sub>-CO-(CH<sub>2</sub>)<sub>n'</sub>-aryl, -(CH<sub>2</sub>)<sub>n</sub>-CO-(CH<sub>2</sub>)<sub>n'</sub>-heterocyclyl, -(CH<sub>2</sub>)<sub>n</sub>-CONH-alkyl, -(CH<sub>2</sub>)<sub>n</sub>-CONH-cycloalkyl, -(CH<sub>2</sub>)<sub>n</sub>-CONH-(CH<sub>2</sub>)<sub>n'</sub>-aryl, -(CH<sub>2</sub>)<sub>n</sub>-CONH-(CH<sub>2</sub>)<sub>n'</sub>-heterocyclyl, -(CH<sub>2</sub>)<sub>n</sub>-NHCO-alkyl, -(CH<sub>2</sub>)<sub>n</sub>-NHCO-cycloalkyl, -(CH<sub>2</sub>)<sub>n</sub>-NHCO-(CH<sub>2</sub>)<sub>n'</sub>-aryl, -(CH<sub>2</sub>)<sub>n</sub>-NHCO-(CH<sub>2</sub>)<sub>n'</sub>-heterocyclyl, -(CH<sub>2</sub>)<sub>n</sub>-NHSO<sub>2</sub>-alkyl, -(CH<sub>2</sub>)<sub>n</sub>-NHSO<sub>2</sub>-cycloalkyl, -(CH<sub>2</sub>)<sub>n</sub>-NHSO<sub>2</sub>-(CH<sub>2</sub>)<sub>n'</sub>-aryl or -(CH<sub>2</sub>)<sub>n</sub>-NHSO<sub>2</sub>-(CH<sub>2</sub>)<sub>n'</sub>-heterocyclyl,  
25 wherein the said alkyl, cycloalkyl, aryl and heterocyclyl groups are unsubstituted or substituted by one to three

substituents selected from chloro, fluoro, hydroxy, methoxy, and C<sub>1</sub>-C<sub>4</sub> alkyl; and n and n', being the same or different,  
30 are zero or an integer from 1 to 5, and R<sub>4</sub>, R<sub>5</sub> are as defined in claim 1.



4. A compound according to claim 2 wherein W, R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are as defined in claim 1;

R<sub>4</sub> is a group of formula (C):



5 wherein R<sub>6</sub> is phenyl, pyridyl, isopropyl, sec-butyl, tert-butyl, adamantyl, cyclopentyl, cyclohexyl, cyclohexylmethyl, phenylmethyl, -pyridylmethyl, 3-pyridylmethyl, 4-pyridylmethyl, 2-thienylmethyl, 3-indolylmethyl, 3-(N-methyl)indolylmethyl, 1,1-diphenylmethyl, 1-naphthyl, 2-naphthyl, 1-naphthylmethyl, 2-naphthylmethyl, quinolylmethyl or isoquinolylmethyl, or a derivative thereof substituted by one to three substituents selected from chloro, fluoro, hydroxy, methoxy, carbamoyl and C<sub>1</sub>-C<sub>4</sub> alkyl; or R<sub>6</sub> is -

10 C(CH<sub>3</sub>)<sub>2</sub>SCH<sub>3</sub> or a sulfoxide or sulfone thereof, -C(CH<sub>3</sub>)<sub>2</sub>OCH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-OCH<sub>3</sub>, or -CH(CH<sub>3</sub>)OH or the tert-butyl ether thereof; and wherein R<sub>7</sub> is -NH<sub>2</sub> or a group -NH-A-, NH-CH<sub>2</sub>-A or NH-CH<sub>2</sub>CH<sub>2</sub>-A, wherein A, being as defined above, is selected from


20 methyl, isopropyl, tert-butyl, phenyl, 4-methoxyphenyl, 4-fluorophenyl, 4-(aminosulfonyl)phenyl, 4-(dimethylamino-sulfonylmethyl)phenyl, (3,4-methylenedioxy)phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 4-piperidyl, 2-thiazolyl, 1-naphthyl, 2-naphthyl, 3-quinolyl, 5-quinolyl, 3-isoquinolyl, 5-isoquinolyl, 3-quinuclidinyl, 2-benzimidazolyl, 5-tetrazolyl,

25 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 1-morpholino, 1-piperidino, or 1-pyrrolidino; and R<sub>5</sub> is hydrogen or methyl.

120

5. A compound according to claim 2 wherein:

wherein W, R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are as defined in claim 1, R<sub>4</sub> is a group of formula (D):




5 wherein R<sub>8</sub> is methyl, ethyl, phenyl, pyridyl, isopropyl, sec-butyl, tert-butyl, adamantyl, cyclopentyl, cyclohexyl, cyclohexylmethyl, phenylmethyl, 3,4-(methylenedioxy)phenyl, piperonyl, pyridylmethyl, thienylmethyl, 3-indolylmethyl, 3-(N-methyl)indolylmethyl, 1,1-diphenylmethyl, naphthyl, naphthylmethyl, quinolylmethyl, isoquinolylmethyl, thiazolyl, thiazolylmethyl, imidazolyl, imidazolylmethyl, or a derivative thereof substituted by one to three substituents selected from chloro, fluoro, hydroxy, methoxy, and C<sub>1</sub>-C<sub>4</sub> alkyl; or R<sub>8</sub> is -C(CH<sub>3</sub>)<sub>2</sub>SCH<sub>3</sub> or a sulfoxide or sulfone

10 thereof, -C(CH<sub>3</sub>)<sub>2</sub>OCH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-OCH<sub>3</sub>, or -CH(CH<sub>3</sub>)OH or a tert-butyl ether thereof; and R<sub>9</sub> is either hydrogen or a group selected from methyl, ethyl, phenyl, 3,4-(methylenedioxy)phenyl, piperonyl, pyridyl, benzimidazolyl and 4-tetrazolyl, which group is unsubstituted or substituted

20 by one to three substituents selected from chloro, fluoro, hydroxy, methoxy, methoxycarbonyl, ethoxycarbonyl and C<sub>1</sub>-C<sub>4</sub> alkyl; and R<sub>5</sub> is hydrogen or methyl group.

6. A compound according to claim 2 wherein:

25 W, R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are as defined in claim 1, R<sub>4</sub> is a 3,4-dihydrocarbostyryl derivative of formula (D'):



(D')

wherein the nitrogen atom may be substituted by methyl, ethyl, propyl, methoxycarbonyl or ethoxycarbonyl, and the phenyl ring may be substituted by one or two substituents  
 5 selected from chloro, fluoro, methyl, methoxy or 3,4-methylenedioxy; and

R<sub>5</sub> is hydrogen or methyl group.

7. A compound according to claim 2 wherein W, R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are as defined in claim 1, R<sub>4</sub> and R<sub>5</sub>, taken together with the  
 10 nitrogen atom to which they are attached, constitute an azaheterocyclic ring selected from morpholine, thiomorpholine, pyrrolidine, piperidine, piperazine, pyridazine, thiazolidine, tetrahydroisoquinoline, hexamethyleneimine and hexahydropyridazine, either  
 15 unsubstituted or substituted by one or more substituents selected from methyl, ethyl, phenyl, 4-fluorophenyl, benzyl, alpha-methylbenzyl, hydroxy, hydroxymethyl, and carbamoyl, or by a group -CONH-A wherein A is selected from methyl, isopropyl, tert-butyl, cyclopropyl cyclobutyl, cyclopentyl,  
 20 cyclohexyl, phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 4-piperidyl, 2-thiazolyl, 1-naphthyl, 2-naphthyl, 3-quinolyl, 5-quinolyl, 3-isoquinolinyl, 5-isoquinolyl, 3-quinuclidinyl, 3,4-(methylenedioxy)phenyl and piperonyl.

25 8. A compound according to claim 2 wherein:

W is either -NHOH or OH;

R<sub>2</sub> is hydroxy;

R<sub>3</sub> is as defined in claim 3;

R<sub>4</sub> and R<sub>5</sub> are as defined in any of claim 4-7;

R<sub>1</sub> is a hydroxymethyl derivative which is -CH<sub>2</sub>OCH<sub>3</sub>, -CH<sub>2</sub>-O-(tetrahydropyranyl), -CH<sub>2</sub>OCOCH<sub>3</sub>, -CH<sub>2</sub>OCOC(CH<sub>3</sub>)<sub>3</sub>, -CH<sub>2</sub>OCONH<sub>2</sub>, -  
5 CH<sub>2</sub>OCONHCH<sub>3</sub>, -CH<sub>2</sub>OCON(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>OCONH-cyclohexyl, -CH<sub>2</sub>OCON(CH<sub>3</sub>)-cyclohexyl, -CH<sub>2</sub>-O-CO-morpholino, -CH<sub>2</sub>-O-CO-pyrrolidino, -CH<sub>2</sub>-O-CO-piperidino; or it is -CH<sub>2</sub>OCH<sub>2</sub>Ph, -CH<sub>2</sub>OCOPh, -CH<sub>2</sub>OCOCH<sub>2</sub>Ph, -CH<sub>2</sub>OCONHCH<sub>2</sub>Ph or a derivative thereof wherein the Ph group is substituted by one, two or three  
10 substituents selected from chloro, fluoro, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, methyl and carbamoyl, or by 3,4-dioxymethylene.

9. A compound according to claim 2 wherein:

W is either -NHOH or OH;

15 R<sub>2</sub> is hydroxy;

R<sub>3</sub> is as defined in claim 3;

R<sub>4</sub> and R<sub>5</sub> are as defined in any of claim 4-7;

R<sub>1</sub> is a mercaptomethyl derivative which is a group -CH<sub>2</sub>S-COCH<sub>3</sub>, -CH<sub>2</sub>S-COPh, -CH<sub>2</sub>S-CH<sub>3</sub>, -CH<sub>2</sub>S-Ph, -CH<sub>2</sub>S-CH<sub>2</sub>Ph, -CH<sub>2</sub>SO<sub>2</sub>-  
20 CH<sub>3</sub>, -CH<sub>2</sub>SO<sub>2</sub>-Ph, -CH<sub>2</sub>SO<sub>2</sub>-CH<sub>2</sub>Ph, -CH<sub>2</sub>S-thienyl, -CH<sub>2</sub>S-(1-methyl-1,2,3,4-tetrazol-5-yl), or a derivative thereof wherein the Ph group is substituted by one or two substituents selected from chloro, fluoro, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, methyl and carbamoyl, or by 3,4-dioxymethylene.

25 10. A compound according to claim 2 wherein:

W is either -NHOH or OH;

R<sub>1</sub> is either hydroxymethyl or mercaptomethyl, or it is a hydroxymethyl derivative as defined in claim 7, or it is a mercaptomethyl derivative as defined in claim 8;


30 R<sub>2</sub> is hydroxy;

123

$R_3$  is isobutyl, cyclopentylmethyl, 3-(4-chlorophenyl)propyl, 3-(4-methoxyphenyl)propyl, 3-(4-biphenyl)propyl, 3-[4-(4-fluorophenyl)phenyl]propyl, 3-[4-(4-chlorophenyl)phenyl]propyl, 3-[4-(4-methoxyphenyl)phenyl]propyl, 3-(4-phenoxyphenyl)propyl, 3-(4-pyridoxyphenyl)propyl, (4-methoxybenzene)sulfonylmethyl, and (4-butoxybenzene)sulfonylmethyl; and  $R_4$  and  $R_5$  are as defined in any of claim 4-7.

11. A process for producing a compound as defined in claim 1,  
 15 which process comprises:


(a) coupling an acid of formula (II), or a salt thereof:



wherein  $R_1$ ,  $R_2$  and  $R_3$  are as defined in claim 1 and  $W'$  is  
 15 protected -OH or protected -NHOH, with an amine of formula (III) or a salt thereof:




wherein  $R_4$  and  $R_5$  are as defined in claim 1, to obtain a compound of formula (Ia):



wherein  $W'$ ,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are as defined above, and then removing the protecting group from this compound to yield a compound of formula (I) wherein  $W$  is  $-OH$  or  $-NHOH$  or a salt thereof; or


- 5 (b) converting a compound of formula (IV)



- wherein  $W''$  is  $OH$  or  $W'$ , and  $R_3$ ,  $R_4$ ,  $R_5$  and  $W'$  are as defined above, into a compound of formula (I) or (Ia) as defined above, and then, when a compound of formula (Ia) is obtained, removing the protecting group from this compound to yield a compound of formula (I) wherein  $W$  is  $-OH$  or  $-NHOH$ , or a salt thereof; or
- 10


- (c) condensing a lactone of formula (V)

15




- wherein  $W'$ ,  $R_1$  and  $R_3$  are as defined above, with an amine of formula (III) as defined above, to obtain a compound of formula (Ia) as defined above wherein  $R_2$  is hydroxy, and then removing the protecting group from this compound to yield a compound of formula (I) wherein  $W$  is  $-OH$  or  $-NHOH$  and  $R_2$  is hydroxy; or
- 20

- (d) condensing a lactone of formula (VI):




- wherein  $W'$ ,  $R_2$  and  $R_3$  are as defined above, with an amine of formula (III) as defined above, to obtain a compound of formula (Ia) above wherein  $R_1$  is hydroxymethyl, and then removing the protecting group from this compound to yield a compound of formula (I) wherein  $W$  is  $-OH$  or  $-NHOH$  and  $R_1$  is hydroxymethyl, and, if desired,
- (e) converting one compound of formula (I) or (Ia) into another compound of formula (I) or (Ia) and/or converting a compound of formula (I) into a pharmaceutically acceptable salt thereof, and/or converting a salt of a compound of formula (I) into the free compound.
12. A pharmaceutical composition which comprises, as active ingredient, a compound as defined in any one of claims 1 to 5 and a pharmaceutically acceptable carrier, diluent or excipient.
13. A compound as defined in any one of claims 1 to 10 for use in a method of treatment or prophylaxis of a disease mediated in a mammal by a matrix metalloproteinase.
14. A compound according to claim 13 for use in a method of treatment or prophylaxis of a tumoral disease in man, in particular for the control of local spread of established tumors, and for the inhibition of the growth of established or occult metastases, either alone or in combination with cytotoxic or cytostatic drugs, or with inhibitors of angiogenesis.

15. A compound according to claim 13 for use in a method of treatment or prophylaxis of rheumatoid arthritis or osteoarthritis in man, alone or in combination with non-steroidal or steroidal antiinflammatory drugs, or with immunosuppressive drugs.
16. A compound as defined in any one of claims 1 to 10 for use in a method of treatment or prophylaxis of an inflammatory, infectious or immunological disease promoted by local or systemic release of soluble TNF.
- 10 17. An acid of formula (II), or an ester thereof:



wherein W' is protected -OH or protected -NHOH, and R<sub>1</sub> to R<sub>3</sub> are as defined in claim 1.


- 15 18. A propiolactone of formula (V):



wherein W' is either free or protected -OH or -NHOH, and R<sub>1</sub> and R<sub>3</sub> are as defined in claim 1.

- 20 19. A butyrolactone of formula (VI):





(VI)

wherein W' is either free or protected -OH or -NHOH, and R<sub>2</sub> and R<sub>3</sub> are as defined in claim 1.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/00531

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D309/12 C07C259/06 C07D213/75 C07C271/24 C07D257/04  
C07C259/08 C07D303/48 C07D295/18 A61K31/16 A61K31/22  
A61K31/395 //C07C69/675, C07C69/78, C07D307/32, C07D305/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 33166 A (THE DU PONT MERCK PHARMACEUTICAL CO) 24 October 1996 see page 29, scheme 1; page 33, scheme 4; page 42, compounds 32-35; claims 1,3,5-13 ---	1,11-16
A	WO 95 29892 A (THE DU PONT MERCK PHARMACEUTICAL CO) 9 November 1995 see schemes 1, 3, 6; table 1, compounds 5-14, 19-28, 33-42, 47-56, 61-69; table 2, compounds 207-214; table 4, compound 403; pages 113-122 ---	1-3,7-16
A	WO 95 19956 A (BRITISH BIOTECH PHARMACEUTICALS LTD) 27 July 1995 see page 22, 2; pages 23-26; examples 4-6, 8-11, 15-16, 32-36 ---	1-4,9-16
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

3 July 1998

Date of mailing of the international search report

10/07/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Van Amsterdam, L

# INTERNATIONAL SEARCH REPORT

In: ational Application No

PCT/EP 98/00531

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>US 5 594 006 A (M. SAKAMOTO ET AL) 14  January 1997  see column 3, line 42 -column 4, line 4;  schemes 5-9; examples 4-6, 96  -----</p>	<p>1-3,6,  9-15</p>

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/00531

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9633166 A	24-10-1996	US 5691381 A AU 5556396 A EP 0821669 A	25-11-1997 07-11-1996 04-02-1998
WO 9529892 A	09-11-1995	AU 2394795 A	29-11-1995
WO 9519956 A	27-07-1995	AT 165817 T AU 682920 B AU 1459795 A BR 9506535 A CA 2181570 A CN 1138851 A DE 19581347 T DE 69502378 D EP 0740652 A EP 0822186 A FI 962904 A GB 2299334 A,B GB 2316078 A,B HU 75059 A JP 9508361 T NO 963030 A NZ 278627 A PL 315745 A SK 94196 A US 5747514 A ZA 9500480 A	15-05-1998 23-10-1997 08-08-1995 16-09-1997 27-07-1995 25-12-1996 05-12-1996 10-06-1998 06-11-1996 04-02-1998 19-07-1996 02-10-1996 18-02-1998 28-03-1997 26-08-1997 19-09-1996 24-04-1997 25-11-1996 05-03-1997 05-05-1998 07-02-1996
US 5594006 A	14-01-1997	JP 8081443 A AU 672888 B AU 6263794 A CA 2136108 A CN 1105799 A EP 0641323 A JP 7157470 A WO 9421612 A	26-03-1996 17-10-1996 11-10-1994 29-09-1994 26-07-1995 08-03-1995 20-06-1995 29-09-1994